

A MILNOR-MOORE TYPE THEOREM FOR PRIMITIVELY GENERATED BRAIDED BIALGEBRAS

ALESSANDRO ARDIZZONI

ABSTRACT. A braided bialgebra is called primitively generated if it is generated as an algebra by its space of primitive elements. We prove that any primitively generated braided bialgebra is isomorphic to the universal enveloping algebra of its infinitesimal braided Lie algebra, notions hereby introduced. This result can be regarded as a Milnor-Moore type theorem for primitively generated braided bialgebras and leads to the introduction of a concept of braided Lie algebra for an arbitrary braided vector space.

CONTENTS

1. Introduction	1
2. Preliminaries	4
3. Pre-categorical and categorical subspaces	6
4. Braided Ideals	8
5. Universal enveloping algebra	11
6. The main result	15
7. The trivial bracket	22
References	24

1. INTRODUCTION

The main Idea. Let K be a fixed field. All vector spaces are understood as vector spaces over K . Let H be a pointed Hopf algebra (i.e. all its simple subcoalgebras are one-dimensional) and denote by G the set of grouplike elements in H . Let $\text{gr}H$ be the graded coalgebra associated to the coradical filtration of H . It is well known that $\text{gr}H$ is isomorphic, as a Hopf algebra, to the Radford-Majid bosonization $R\#KG$ of R by the group algebra KG , where R , the so-called diagram of H , is a suitable connected braided bialgebra in the braided monoidal category of Yetter-Drinfeld modules over KG . This is the starting point of the *lifting method* for the classification of finite dimensional pointed Hopf algebras, introduced by N. Andruskiewitsch and H.-J. Schneider, see e.g. [AS1]. Accordingly to this method, first one has to describe R by generators and relations, then to lift the informations obtained to H . It is worth to notice that Andruskiewitsch and Schneider conjectured that, in the finite dimensional case and characteristic zero, R is always primitively generated, [AS2, Conjecture 1.4] and [AS3, Conjecture 2.7].

Therefore, in many cases, for proving certain properties of Hopf algebras, one can reduce to the connected case. The price to pay is to work with bialgebras in a braided category, and not with ordinary bialgebras.

Actually in our case it is more convenient to work with braided bialgebras, that were introduced in [Ta]. Recall that a *braided vector space* (V, c) consists of a vector space V and K -linear map $c : V \otimes V \rightarrow V \otimes V$, called braiding, satisfying $c_1 c_2 c_1 = c_2 c_1 c_2$ (the so-called quantum Yang-Baxter equation). Here $c_1 = c \otimes V$ and $c_2 = V \otimes c$. A *braided bialgebra* is then a braided vector space

2000 *Mathematics Subject Classification.* Primary 16W30; Secondary 16S30.

Key words and phrases. Braided bialgebras, braided Lie algebras, universal enveloping algebras.

This paper was written while the author was member of GNSAGA with partial financial support from MIUR within the National Research Project PRIN 2007.

which is both an algebra and a coalgebra with structures suitably compatible with the braiding. Examples of braided bialgebras are all bialgebras in those braided monoidal categories which are monoidal subcategories of the category of vector spaces.

A coalgebra (a fortiori a braided bialgebra) is called connected if its coradical (i.e. the sum of all its simple subcoalgebras) is one-dimensional. The main result of this paper, Theorem 6.9, states that every primitively generated (whence connected) braided bialgebra is isomorphic, as a braided bialgebra, to the universal enveloping algebra of its infinitesimal braided Lie algebra. This result can be seen as an extension of the celebrated Milnor-Moore Theorem [MM, Theorem 5.18] for cocommutative connected bialgebras (once observed that such a bialgebra is always primitively generated): in characteristic zero, any cocommutative connected bialgebra is the enveloping algebra of its space of primitive elements, regarded as a Lie algebra in a canonical way.

In order to understand the statement of the main theorem, we need to clarify what is meant by infinitesimal braided Lie algebra of a braided bialgebra and by universal enveloping algebra of it. The definition of infinitesimal braided Lie algebra requires that of the braided tensor algebra $T(V, c)$ associated to a braided vector space (V, c) . As an algebra, $T(V, c)$ is the tensor algebra $T = T(V)$. Then T can be endowed with a bialgebra structure depending on c which is denoted by $T(V, c)$ and called the braided tensor algebra of (V, c) .

Now, let A be a braided bialgebra. The braiding of A induces a braiding c_P on the space $P = P(A)$ of primitive elements in A . The braided vector space (P, c_P) will be called the infinitesimal part of A .

Let us construct a direct system of braided bialgebras

$$U^{[0]} \xrightarrow{\pi_0^1} U^{[1]} \xrightarrow{\pi_1^2} U^{[2]} \xrightarrow{\pi_2^3} \dots$$

The first term is $U^{[0]} := T(P, c_P)$. The usual universal property of $T(P)$ yields an algebra map $\varphi^{[0]} : U^{[0]} \rightarrow A$, which turns out to be a braided bialgebra map, such that the diagram

$$\begin{array}{ccc} P & \xrightarrow{i_{U^{[0]}}} & U^{[0]} \\ & \searrow i & \swarrow \varphi^{[0]} \\ & A & \end{array}$$

commutes, where $i_{U^{[0]}}$ is the canonical map and i is the inclusion. Set $P^{[0]} := P(U^{[0]})$. Then $i_{U^{[0]}}$ corestricts to $i^{[0]} : P \rightarrow P^{[0]}$. Moreover, $\varphi^{[0]}$ preserves primitive elements so that it induces a map $b^{[0]} : P^{[0]} \rightarrow P$. The second term $U^{[1]}$ of the direct system is then defined as the quotient of $U^{[0]}$ by its two-sided ideal generated by $[\text{Id} - i^{[0]}b^{[0]}] [P^{[0]}]$. From the definition one gets $\varphi^{[0]}i^{[0]}b^{[0]} = \varphi^{[0]}$ so that $\varphi^{[0]}$ quotients to a bialgebra homomorphism $\varphi^{[1]} : U^{[1]} \rightarrow A$ such that the diagram

$$\begin{array}{ccc} U^{[0]} & \xrightarrow{\pi_0^1} & U^{[1]} \\ & \searrow \varphi^{[0]} & \swarrow \varphi^{[1]} \\ & A & \end{array}$$

commutes, where π_0^1 denotes the canonical projection. Set $P^{[1]} := P(U^{[1]})$. Then $i_{U^{[1]}} = \pi_0^1 i_{U^{[0]}}$ corestricts to $i^{[1]} : P \rightarrow P^{[1]}$. Moreover, $\varphi^{[1]}$ induces a map $b^{[1]} : P^{[1]} \rightarrow P$. The third term $U^{[2]}$ of the direct system is then the quotient of $U^{[1]}$ by its two-sided ideal generated by $(\text{Id} - i^{[1]}b^{[1]}) [P^{[1]}]$. Moreover $\varphi^{[1]}$ quotients to a bialgebra homomorphism $\varphi^{[2]}$ such that $\varphi^{[2]}\pi_1^2 = \varphi^{[1]}$.

Going on this way, we obtain the direct system $(U^{[n]})_{n \in \mathbb{N}}$ and a sequence of maps $b_P := (b^{[n]})_{n \in \mathbb{N}}$, where $b^{[n]} : P^{[n]} \rightarrow P$. The datum (P, c_P, b_P) is called the infinitesimal braided Lie algebra of A .

The direct limit of the direct system $(U^{[n]})_{n \in \mathbb{N}}$ is a braided bialgebra that we denote by

$$U(P, c_P, b_P) := \varinjlim U^{[n]}.$$

The compatible family of morphisms $(\varphi^{[n]})_{n \in \mathbb{N}}$ gives rise to a braided bialgebra homomorphism $\varphi^{[\infty]} : U(P, c_P, b_P) \rightarrow A$. We can show that the space of primitive elements in $U(P, c_P, b_P)$ identifies with P . Using this fact, in Theorem 6.9, by applying a famous result due to Heyneman

and Radford (see [Mo, Theorem 5.3.1]), we can prove that $\varphi^{[\infty]}$ is an isomorphism whenever A is primitively generated. Now, the structure and properties of $U(P, c_P, b_P)$ are encoded in the datum (P, c_P, b_P) . Indeed this leads to the introduction of what will be called a braided Lie algebra (V, c, b) for any braided vector space (V, c) and of the related universal enveloping algebra $U(V, c, b)$.

The results in Detail. The paper is organized as follows.

Section 2 contains preliminary facts and notations that will be used in the paper.

Section 3 concerns general results on pre-categorical and categorical spaces that will be applied in next section.

Section 4 deals with the problem of quotienting braided bialgebras. In particular, we investigate braided bialgebra quotients of the braided tensor algebra $T(V, c)$ associated to a given braided vector space (V, c) . In Lemma 4.10, these braided bialgebras are characterized as (V, c) -bialgebras, a notion hereby introduced. In Theorem 4.11, we show how to construct a quotient of a (V, c) -bialgebra whenever there exists a suitable map. This fact will be relevant in defining a braided Lie algebra of a braided vector space.

Section 5, is devoted to the introduction and investigation of the notion of braided Lie algebra and universal enveloping algebra. Let (V, c) be a braided vector space. We say that (V, c) has a bracket if there is a family of maps $b = (b^{[n]} : P^{[n]} \rightarrow V)_{n \in \mathbb{N}}$ which allow, roughly speaking, to construct step by step a direct system of braided bialgebras

$$T(V, c) = U^{[0]} \xrightarrow{\pi_0^1} U^{[1]} \xrightarrow{\pi_1^2} U^{[2]} \xrightarrow{\pi_2^3} \dots$$

as we did in the case of (P, c_P) . The direct limit of this direct system is a braided bialgebra denoted by

$$U(V, c, b) := \varinjlim U^{[n]}.$$

This bialgebra is called the universal enveloping algebra of (V, c, b) . We say that (V, c, b) is a *braided Lie algebra* whenever (V, c) is a braided vector space, b is a bracket on (V, c) and the canonical map $i_U : V \rightarrow U(V, c, b)$ is injective (the latter will be called the *implicit Jacobi identity*). In Corollary 5.6, we prove that, for a braided Lie algebra (V, c, b) , the space of primitive elements in $U(V, c, b)$ identifies with V through the canonical map $i_U : V \rightarrow U$. In Proposition 5.7, we get that for a braided vector space (V, c) with a bracket b , the implicit Jacobi identity holds if and only if $b^{[n]}i^{[n]} = \text{Id}$ for all $n \in \mathbb{N}$. Applying this result, in Corollary 5.8, we give the alternative description of $U^{[n+1]}$ as the quotient of $U^{[n]}$ by its two-sided ideal generated by $\text{Ker}(b^{[n]})$.

Let (V, c, b) be a braided Lie algebra and let $U = U(V, c, b)$. In Theorem 5.10, we associate to (V, c, b) a braided Lie algebra (V, c, β) in the sense of [Ar2]. Furthermore, we show that the corresponding universal enveloping algebra $\mathbb{U}(V, c, \beta)$, also defined in [Ar2], coincide with $U^{[1]}$ as given above. Therefore, the notions defined in [Ar2] can be seen as a first step of the procedure we consider in the present paper. Now, for a given braided vector space (V, c) , let $E(V, c)$ be the vector space spanned by primitive elements in $T(V, c)$ of homogeneous degree at least two. Let $\mathcal{S}$ be the class of those braided vector spaces (V, c) whose Nichols algebra $\mathcal{B}(V, c)$ is obtained dividing out $T(V, c)$ by its two-sided ideal generated by $E(V, c)$. In Theorem 5.10, we also prove that $U(V, c, b) = \mathbb{U}(V, c, \beta)$ whenever (V, c) is in $\mathcal{S}$ (therefore, the main theorem of the present paper extends [Ar2, Theorem 5.7]). It is remarkable that, see [Ar1], this happens, for instance, when

- $c^2 = \text{Id}$ and $\text{char} K = 0$;
- there exists $q \in K$, not a root of unity, such that $(c + \text{Id}_{V^{\otimes 2}}) \circ (c - q\text{Id}_{V^{\otimes 2}}) = 0$ i.e. c is of Hecke type with mark q ;
- the Nichols algebra $\mathcal{B}(V, c)$ is quadratic.
- (V, c) is a braided vector space of diagonal type such that $\mathcal{B}(V, c)$ is a domain and its Gelfand-Kirillov dimension is finite.

Section 6 encloses the main results of the paper. In Proposition 6.1, we prove that any braided algebra A becomes a braided Lie algebra in a canonical way. In Theorem 6.5 we show how this Lie algebra structure restricts to the space $P = P(A)$ of primitive elements in A when the braided algebra is further a braided bialgebra. This braided Lie algebra on P is called the infinitesimal braided Lie algebra of A (it is constructed as explained in the first part of the introduction). In

Theorem 6.7 a universal property for the universal enveloping algebra of a braided Lie algebra is obtained. Theorems 6.8 and 6.9, constitutes together a Milnor-Moore type theorem for primitively generated braided bialgebras. Explicitly, Theorem 6.8 establish that the braided Lie algebra (V, c, b) identifies with the infinitesimal braided Lie algebra of $U(V, c, b)$. Theorem 6.9, which is the main result of the paper, states that every primitively generated braided bialgebra is isomorphic as a braided bialgebra to the universal enveloping algebra of its infinitesimal braided Lie algebra. In Example 6.10, we show how to recover the ordinary notion of Lie algebra from that of braided Lie algebra.

Section 7, regards the concept of trivial bracket. In Example 7.1, we associates the so-called trivial bracket b_{tr} to any braided vector space (V, c) . This way (V, c, b_{tr}) becomes a braided Lie algebra and $U(V, c, b_{tr})$ is nothing but the Nichols algebra $\mathcal{B}(V, c)$ associated to (V, c) . In Proposition 7.2, we establish a condition which guarantees that the bracket of a braided Lie algebra is trivial. In Example 7.3, we apply this result to give an example of a braided vector space (V, c) which becomes a braided Lie algebra only through the trivial bracket. It is remarkable that this braided vector space does not belong to the class $\mathcal{S}$ mentioned above.

2. PRELIMINARIES

Throughout this paper K will denote a field. All vector spaces will be defined over K and the tensor product over K will be denoted by $\otimes$.

In this section we define the main notions that we will deal with in the paper.

Definition 2.1. Let V be a vector space over a field K . A K -linear map $c = c_V : V \otimes V \rightarrow V \otimes V$ is called a **braiding** if it satisfies the quantum Yang-Baxter equation

$$(2.1) \quad c_1 c_2 c_1 = c_2 c_1 c_2$$

on $V \otimes V \otimes V$, where we set $c_1 := c \otimes V$ and $c_2 := V \otimes c$. The pair (V, c) will be called a **braided vector space**. A morphism of braided vector spaces (V, c_V) and (W, c_W) is a K -linear map $f : V \rightarrow W$ such that $c_W(f \otimes f) = (f \otimes f)c_V$.

A general method for producing braided vector spaces is to take an arbitrary braided category $\mathcal{M}$ which is a monoidal subcategory of the category of K -vector spaces. Hence any object $V \in \mathcal{M}$ can be regarded as a braided vector space with respect to $c := c_{V, V}$, where $c_{X, Y} : X \otimes Y \rightarrow Y \otimes X$ denotes the braiding in $\mathcal{M}$, for all $X, Y \in \mathcal{M}$. Both the category of comodules over a coquasitriangular Hopf algebra and the category of Yetter-Drinfeld modules over a Hopf algebra with bijective antipode are examples of such categories. More particularly, every bicharacter of a group G induces a braiding on the category of G -graded vector spaces.

Definition 2.2. [Ba] A quadruple (A, m, u, c) is called a **braided algebra** if

- (A, m, u) is an associative unital algebra;
- (A, c) is a braided vector space;
- m and u commute with c , that is the following conditions hold:

$$(2.2) \quad c(m \otimes A) = (A \otimes m)(c \otimes A)(A \otimes c),$$

$$(2.3) \quad c(A \otimes m) = (m \otimes A)(A \otimes c)(c \otimes A),$$

$$(2.4) \quad c(u \otimes A) = A \otimes u, \quad c(A \otimes u) = u \otimes A.$$

A morphism of braided algebras is, by definition, a morphism of ordinary algebras which, in addition, is a morphism of braided vector spaces.

A quadruple $(C, \Delta, \varepsilon, c)$ is called a **braided coalgebra** if

- (C, Δ, ε) is a coassociative counital coalgebra;
- (C, c) is a braided vector space;
- Δ and ε commute with c , that is the following relations hold:

$$(2.5) \quad (\Delta \otimes C)c = (C \otimes c)(c \otimes C)(C \otimes \Delta),$$

$$(2.6) \quad (C \otimes \Delta)c = (c \otimes C)(C \otimes c)(\Delta \otimes C),$$

$$c_T^{n,m} = \begin{array}{c} \begin{array}{cc} T^n(V) & T^m(V) \end{array} \\ \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} \\ \begin{array}{cc} T^m(V) & T^n(V) \end{array} \end{array}$$

$$(2.7) \quad (\varepsilon \otimes C)c = C \otimes \varepsilon, \quad (C \otimes \varepsilon)c = \varepsilon \otimes C.$$

A morphism of braided coalgebras is, by definition, a morphism of ordinary coalgebras which, in addition, is a morphism of braided vector spaces.

[Ta, Definition 5.1] A sextuple $(B, m, u, \Delta, \varepsilon, c)$ is called a **braided bialgebra** if

- (B, m, u, c) is a braided algebra
- $(B, \Delta, \varepsilon, c)$ is a braided coalgebra
- the following relations hold:

$$(2.8) \quad \Delta m = (m \otimes m)(B \otimes c \otimes B)(\Delta \otimes \Delta).$$

Examples of the notions above are algebras, coalgebras and bialgebras in any braided category $\mathcal{M}$ which is a monoidal subcategory of the category of K -vector spaces. The notion of braided bialgebra admits a graded counterpart which is called graded braided bialgebra. For further results on this topic the reader is referred to [Ar1].

Example 2.3. Let (V, c) be a braided vector space. Consider the tensor algebra $T = T(V)$ and let m_T and u_T denote its multiplication and unit respectively. This is a graded braided algebra with n -th graded component $T^n(V) = V^{\otimes n}$. The braiding c_T on T is defined using the braiding of V :

Now $T \otimes T$ becomes itself an algebra with multiplication $m_{T \otimes T} := (m_T \otimes m_T) \circ (T \otimes c_T \otimes T)$. This algebra is denoted by $T \otimes_c T$. The universal property of the tensor algebra yields two algebra homomorphisms $\Delta_T : T \rightarrow T \otimes_c T$ and $\varepsilon_T : T \rightarrow K$. It is straightforward to check that $(T, m_T, u_T, \Delta_T, \varepsilon_T, c_T)$ is a graded braided bialgebra.

Definition 2.4. The graded braided bialgebra described in Example 2.3 is called the **braided tensor algebra** and is denoted by $T(V, c)$.

Note that Δ_T really depends on c . For instance, one has $\Delta_T(z) = z \otimes 1 + 1 \otimes z + (c + \text{Id})(z)$, for all $z \in V \otimes V$.

2.5. Recall that a coalgebra C is called **connected** if the coradical C_0 of C (i.e the sum of all simple subcoalgebras of C) is one dimensional. In this case there is a unique group-like element $1_C \in C$ such that $C_0 = K1_C$. A morphism of connected coalgebras is just a coalgebra homomorphism (clearly it preserves the grouplike element).

By definition, a braided coalgebra (C, c) is **connected** if the underlying coalgebra is connected and, for any $x \in C$, $c(x \otimes 1_C) = 1_C \otimes x$ and $c(1_C \otimes x) = x \otimes 1_C$.

Given a coalgebra C with a distinguished grouplike element 1_C , we will use the following notation

$$\delta = \delta_C : C \rightarrow C \otimes C, \quad \delta(c) := c \otimes 1_C + 1_C \otimes c - \Delta(c) \quad \text{for all } c \in C.$$

Remark 2.6. Let $C = \bigoplus_{n \in \mathbb{N}} C^n$ be a graded braided coalgebra. By [Sw, Proposition 11.1.1], if $(C_n)_{n \in \mathbb{N}}$ is the coradical filtration, then $C_n \subseteq \bigoplus_{m \leq n} C^m$. Therefore, C is connected if C^0 is a one dimensional vector space.

Definition 2.7. A braided bialgebra B is called **primitively generated** if it is generated as an algebra by the space $P(B) = \text{Ker} \delta_B = \{b \in B \mid \Delta(b) = b \otimes 1_B + 1_B \otimes b\}$ of primitive elements in B . See [MM, page 239].

Example 2.8. Let B be a connected ordinary bialgebra in characteristic zero.

Assume that B is primitively generated and set $P = P(A)$. Then, we have a Hopf algebra projection $T(P) \rightarrow B$. Since $T(P)$ is cocommutative, B is cocommutative too.

Conversely, if B is cocommutative, then $\text{gr}B$ is also cocommutative connected. By [AMS, (3) $\Rightarrow$ (4) in Theorem 2.15], we have that $\text{gr}B$ is strictly generated as an algebra by $\text{gr}^1 B = B_1/B_0$. Hence B is primitively generated. Note that we did not use Milnor-Moore theorem to conclude that B is primitively generated.

Proposition 2.9. [Ar2, Proposition 5.8] *Let B be a primitively generated braided bialgebra. Then the underlying braided coalgebra is connected.*

Example 2.10. We give an example of a finite dimensional connected braided bialgebra which is not primitively generated (it is a particular case of the graded dual of S as in [AS3, Example 2.5]). Assume $\text{char}K = 2$. Consider the algebra B generated as an algebra over K by x and y with relations $x^2, y^2, yx - xy$. Then $1, x, y, xy$ form a basis for B . We can regard B as an ordinary bialgebra by setting

$$\Delta_B(x) = 1 \otimes x + x \otimes 1, \quad \Delta_B(y) = 1 \otimes y + x \otimes x + y \otimes 1.$$

Clearly B is not primitively generated. On the other hand, if we set $B(0) := K, B(1) = K + Kx, B(2) := K + Kx + Ky$, and $B(3) = B$, we get a filtration for the coalgebra B . By [Sw, Proposition 11.1.1], this yields $\text{Corad}(B) \subseteq B(0) = K$. Hence B is connected.

Remark 2.11. It is remarkable that, in characteristic zero, in the literature there is no example of a finite dimensional connected braided bialgebra which is not primitively generated. See also [AS2, Conjecture 1.4] and [AS3, Conjecture 2.7].

Lemma 2.12. *Let $((B_u)_{u \in \mathbb{N}}, (\xi_u^v)_{u,v \in \mathbb{N}})$ be a direct system of vector spaces, where, for $u \leq v$, $\xi_u^v : B_u \rightarrow B_v$. Assume that each B_u is endowed with a braided bialgebra structure such that ξ_u^v is a braided bialgebra homomorphism, for every $u, v \in \mathbb{N}$. Then $\varinjlim B_u$ carries a natural braided bialgebra structure that makes it the direct limit of $((B_u)_{u \in \mathbb{N}}, (\xi_u^v)_{u,v \in \mathbb{N}})$ as a direct system of braided bialgebras.*

Proof. It is straightforward. □

3. PRE-CATEGORICAL AND CATEGORICAL SUBSPACES

In this section we investigate the properties of pre-categorical and categorical vector spaces that will be needed next section.

Definition 3.1. (cf. [Kh2, 2.2]) A subspace L of a braided vector space (V, c) will be called **pre-categorical** whenever

$$(3.1) \quad c(L \otimes V + V \otimes L) \subseteq L \otimes V + V \otimes L.$$

Proposition 3.2. *Let (V, c) be a braided vector space.*

1) *Let $f : (V, c) \rightarrow (V', c')$ be a morphism of braided vector spaces. Then, $\text{Ker}(f)$ is a pre-categorical subspace of (V, c) .*

2) *Let L be pre-categorical subspace of (V, c) . Then, the quotient V/L carries a unique braided vector space structure such that the canonical projection is a morphism of braided vector spaces.*

Proof. 1) We have

$$(f \otimes f) c(\text{Ker}(f \otimes f)) = c'(f \otimes f)(\text{Ker}(f \otimes f)) = 0$$

so that $c(\text{Ker}(f \otimes f)) \subseteq \text{Ker}(f \otimes f)$. Since $\text{Ker}(f \otimes f) = \text{Ker}(f) \otimes V + V \otimes \text{Ker}(f)$, we get that $\text{Ker}(f)$ is a pre-categorical subspace of (V, c) .

2) Set $V' := V/L$ and denote by $\pi : V \rightarrow V'$ the canonical projection. We have

$$c(\text{Ker}(\pi \otimes \pi)) = c(L \otimes V + V \otimes L) \stackrel{(3.1)}{\subseteq} L \otimes V + V \otimes L = \text{Ker}(\pi \otimes \pi)$$

so that there exists $c' : V' \otimes V' \rightarrow V' \otimes V'$ such that $c'(\pi \otimes \pi) = (\pi \otimes \pi)c$. By this equality and the fact that π is surjective, it is straightforward to prove that (V', c') is a braided vector space. □

Definition 3.3. A subspace L of a braided vector space (V, c) is called **categorical** (cf. [Ta, Section 6]; see also [AN, 2.5]) whenever

$$(3.2) \quad c(L \otimes V) \subseteq V \otimes L \quad \text{and} \quad c(V \otimes L) \subseteq L \otimes V.$$

In this case c induces maps $c_{L,V} : L \otimes V \rightarrow V \otimes L$ and $c_{V,L} : V \otimes L \rightarrow L \otimes V$.

If L and L' are categorical subspaces of (V, c) , we say that $f : L \rightarrow L'$ is a **morphism of categorical subspaces** of (V, c) if f is a K -linear map such that

$$(3.3) \quad c_{L',V}(f \otimes V) = (V \otimes f)c_{L,V} \quad \text{and} \quad c_{V,L'}(V \otimes f) = (f \otimes V)c_{V,L}.$$

Let L and L' be categorical subspaces of (V, c) . Then

$$c(L \otimes L') \subseteq c(L \otimes V) \cap c(V \otimes L') \subseteq (V \otimes L) \cap (L' \otimes V) = L' \otimes L$$

so that c induces a map $c_{L,L'} : L \otimes L' \rightarrow L' \otimes L$. If in addition $L \subseteq L'$, we get that L is a categorical subspace of L' too.

We will sometime use the simplified notation $c_L := c_{L,L}$.

Remark 3.4. Any categorical subspace of a braided vector space is pre-categorical.

In the connected case, the following result appeared in [Kh2, page 4].

Lemma 3.5. [Ar2, Lemma 2.10] *Let (B, c) be a braided bialgebra. Let $P = P(B)$ be the space of primitive elements of B . Then P is a categorical subspace of (B, c) .*

Definition 3.6. With same assumptions and notations as in Lemma 3.5, the braiding c_P of P (which is induced by c) will be called the **infinitesimal braiding of B** while (P, c_P) will be called the **infinitesimal part of B** .

Lemma 3.7. *Let L and L' be categorical subspaces of (V, c) , and let $f : L \rightarrow L'$ be a morphism of categorical subspaces of (V, c) . Then $\text{Ker}(f)$ and $\text{Im}(f)$ are categorical subspaces of (V, c) .*

Proof. Set $K := \text{Ker}(f)$. Then

$$(V \otimes f)c(K \otimes V) = (V \otimes f)c_{L,V}(K \otimes V) \stackrel{(3.3)}{=} c_{L',V}(f \otimes V)(K \otimes V) = 0$$

so that $c(K \otimes V) \subseteq \text{Ker}(V \otimes f) = V \otimes K$. Similarly one gets $c(V \otimes K) \subseteq K \otimes V$ so that K is a categorical subspace of (V, c) . Set $D := \text{Im}(f)$. We have

$$c(D \otimes V) = c_{L',V}(D \otimes V) = c_{L',V}(f \otimes V)(L \otimes V) \stackrel{(3.3)}{=} (V \otimes f)c_{L,V}(L \otimes V) \subseteq V \otimes D$$

so that $c(D \otimes V) \subseteq V \otimes D$. Similarly one gets $c(V \otimes D) \subseteq D \otimes V$ so that D is a categorical subspace of (V, c) . $\square$

Definition 3.8. Let B be a braided bialgebra. Let V be a vector subspace of B . Then we can define recursively the iterated powers of V in B by setting

$$V \cdot B^0 := K \quad \text{and} \quad V \cdot B^n := V \cdot B^{(n-1)} \cdot_B V, \quad \text{for all } n \geq 1.$$

Lemma 3.9. *Let (B, c) be a braided bialgebra. Let V be a categorical subspace of (B, c) . Then $V \cdot B^n$ is a categorical subspace of B for all $n \in \mathbb{N}$.*

Proof. We prove the statement by induction on $n \in \mathbb{N}$. Set $V^n := V \cdot B^n$. For $n = 0$ it follows by (2.4) and for $n = 1$ there is nothing to prove. Let $n \geq 2$ and assume $V \cdot B^{n-1}$ is a categorical subspace of (B, c) . Let m denote the multiplication of B . Then

$$\begin{aligned} c(V^n \otimes B) &\subseteq c(V^{n-1} \cdot_B V \otimes B) \\ &\subseteq c(m \otimes B)(V^{n-1} \otimes V \otimes B) \\ &\stackrel{(2.2)}{\subseteq} (B \otimes m)(c \otimes B)(B \otimes c)(V^{n-1} \otimes V \otimes B) \\ &\subseteq (B \otimes m)(c \otimes B)(V^{n-1} \otimes B \otimes V) \\ &\stackrel{\text{induction hypothesis}}{\subseteq} (B \otimes m)(B \otimes V^{n-1} \otimes V) \subseteq B \otimes V^n. \end{aligned}$$

Similarly, using (2.3) one proves that $c(B \otimes V^n) \subseteq V^n \otimes B$. $\square$

Proposition 3.10. *Let B be a braided bialgebra generated as an algebra by some subspace V . Then, the following assertions are equivalent.*

- (i) V is a braided subspace of B .
- (ii) V is a categorical subspace of B .

Proof. Denote by c the braiding of B .

(ii) $\Rightarrow$ (i) By Definition 3.1, c induces a map $c_V : V \otimes V \rightarrow V \otimes V$. Clearly (V, c_V) is a braided subspace of (B, c) .

(i) $\Rightarrow$ (ii) We have to check that $c(V \otimes B) \subseteq B \otimes V$ and $c(B \otimes V) \subseteq V \otimes B$. We treat the former inclusion, the latter having a similar proof. Since B is generated by V , then $B = \cup V^n$ where $V^n := V^{\cdot B^n}$ for all $n \in \mathbb{N}$. Thus it suffices to check that $c(V \otimes V^n) \subseteq V^n \otimes V$. For $n = 1$, there is nothing to prove as V is a braided subspace of (B, c) . Assume that the formulas are true for $n - 1$ and denote by m the multiplication of B . Then, we have

$$\begin{aligned}
 c(V \otimes V^n) &= c(B \otimes m)(V \otimes V^{n-1} \otimes V) \\
 &\stackrel{(2.3)}{=} (m \otimes B)(B \otimes c)(c \otimes B)(V \otimes V^{n-1} \otimes V) \\
 &\stackrel{\text{induction hypothesis}}{\subseteq} (m \otimes B)(B \otimes c)(V^{n-1} \otimes V \otimes V) \\
 &\subseteq (m \otimes B)(V^{n-1} \otimes V \otimes V) = V^n \otimes V.
 \end{aligned}$$

□

4. BRAIDED IDEALS

In this section we face the problem of quotienting braided bialgebras. In particular, we investigate braided bialgebra quotients of the braided tensor algebra $T(V, c)$ associated to a given braided vector space (V, c) . These braided bialgebras will be characterized as (V, c) -bialgebras.

Definition 4.1. A **braided ideal** of a braided bialgebra B is a pre-categorical subspace of B which is both an ideal and a coideal of B .

Lemma 4.2. *For a given braided ideal I of a braided bialgebra B , the quotient B/I carries a unique braided bialgebra structure such that the canonical projection is a braided bialgebra homomorphism.*

Proof. Set $A := B/I$. Since I is both an ideal and a coideal of B , it is clear that A is both an algebra and a coalgebra in such a way that the projection $\pi : B \rightarrow A$ is both an algebra and a coalgebra map. Since I is a pre-categorical subspace of B , by Proposition 3.2, the quotient $A = B/I$ carries a unique braided vector space structure such that π is a morphism of braided vector spaces. From this fact, since A is a braided bialgebra, π is surjective and it is both an algebra and a coalgebra map, it is straightforward to prove that B is a braided bialgebra in such a way that π is a braided bialgebra map. □

The following result in the case $B = T(V, c)$ for some braided vector space (V, c) appeared in [Kh2, page 7].

Theorem 4.3. *Let B be a braided bialgebra with infinitesimal part (P, c_P) . Let S be a pre-categorical subspace of (P, c_P) . Then the ideal (S) of B generated by S is a braided ideal of B .*

Proof. Let c, m, Δ and ε denote the braiding, the multiplication, the comultiplication and the counit of B respectively. Set $I := (S) = BSB$. We prove that $c(I \otimes B) \subseteq I \otimes B + B \otimes I$. We have

$$\begin{aligned}
 c(I \otimes B) &= c(BSB \otimes B) \\
 &= c(m \otimes B)(B \otimes m \otimes B)(B \otimes S \otimes B \otimes B) \\
 &= (B \otimes m)(B \otimes B \otimes m)(c \otimes B \otimes B)(B \otimes c \otimes B)(B \otimes S \otimes c(B \otimes B)) \\
 &\subseteq (B \otimes m)(B \otimes B \otimes m)(c \otimes B \otimes B)(B \otimes c \otimes B)(B \otimes S \otimes B \otimes B) \\
 &\stackrel{(3.1)}{\subseteq} (B \otimes m)(B \otimes B \otimes m)(c \otimes B \otimes B)(B \otimes S \otimes B \otimes B + B \otimes B \otimes S \otimes B)
 \end{aligned}$$

$$\begin{aligned}
& \stackrel{(3.1)}{\subseteq} (B \otimes m)(B \otimes B \otimes m)(B \otimes S \otimes B \otimes B + S \otimes B \otimes B \otimes B + B \otimes B \otimes S \otimes B) \\
& \subseteq B \otimes SBB + S \otimes BBB + B \otimes BSB \subseteq B \otimes I + I \otimes B.
\end{aligned}$$

Similarly one gets $c(B \otimes I) \subseteq I \otimes B + B \otimes I$ so that I is a pre-categorical subspace of (B, c) . It remains to check that I is a coideal of B . Note that $S \subseteq P$ so that $\Delta(S) \subseteq B \otimes S + S \otimes B$ and $\varepsilon(S) = 0$. Let us check that $\Delta(I) \in I \otimes B + B \otimes I$:

$$\begin{aligned}
& \Delta(I) \\
&= \Delta(SBS) \\
&= \Delta m(m \otimes B)(B \otimes S \otimes B) \\
&\stackrel{(2.8)}{=} (m \otimes m)(B \otimes c \otimes B)(\Delta \otimes \Delta)(m \otimes B)(B \otimes S \otimes B) \\
&\stackrel{(2.8)}{=} \left[\begin{array}{c} (m \otimes m)(B \otimes c \otimes B)(m \otimes m \otimes B \otimes B) \\ (B \otimes c \otimes B \otimes B \otimes B)(\Delta \otimes \Delta \otimes \Delta)(B \otimes S \otimes B) \end{array} \right] \\
&\subseteq \left[\begin{array}{c} (m \otimes m)(B \otimes c \otimes B)(m \otimes m \otimes B \otimes B) \\ (B \otimes c \otimes B \otimes B \otimes B)(B \otimes B \otimes B \otimes S \otimes B \otimes B + B \otimes B \otimes S \otimes B \otimes B \otimes B) \end{array} \right] \\
&\stackrel{(3.1)}{\subseteq} \left[\begin{array}{c} (m \otimes m)(B \otimes c \otimes B)(m \otimes m \otimes B \otimes B) \\ (B \otimes B \otimes B \otimes S \otimes B \otimes B + B \otimes B \otimes S \otimes B \otimes B \otimes B + B \otimes S \otimes B \otimes B \otimes B \otimes B) \end{array} \right] \\
&\subseteq (m \otimes m)(B \otimes c \otimes B)(B \otimes BS \otimes B \otimes B + B \otimes SB \otimes B \otimes B + BS \otimes B \otimes B \otimes B) \\
&\subseteq (m \otimes m)(B \otimes c \otimes B)(B \otimes I \otimes B \otimes B + I \otimes B \otimes B \otimes B) \\
&\subseteq (m \otimes m)(B \otimes I \otimes B \otimes B + B \otimes B \otimes I \otimes B + I \otimes B \otimes B \otimes B) \\
&\subseteq BI \otimes B + B \otimes IB + IB \otimes B \\
&\subseteq I \otimes B + B \otimes I.
\end{aligned}$$

Furthermore $\varepsilon(I) = \varepsilon(BSB) = \varepsilon(B)\varepsilon(S)\varepsilon(B) = 0$. Therefore I is a coideal. $\square$

Corollary 4.4. *Let B be a braided bialgebra with infinitesimal part (P, c_P) . Let S be a categorical subspace of (P, c_P) . Then the ideal (S) of B generated by S is a braided ideal of B .*

Proof. Apply Remark 3.4 and Theorem 4.3. $\square$

Proposition 4.5. *Let B be a braided bialgebra with infinitesimal part P . Let $b : P \rightarrow P$ be a morphism of categorical subspaces of B . Then the two-sided ideal of B generated by $[\text{Id} - b][P]$ is a braided ideal. Thus the quotient*

$$W(B, b) := \frac{B}{([\text{Id} - b][P])}$$

carries a unique braided bialgebra structure such that the canonical projection is a braided bialgebra homomorphism.

Proof. Since both $\text{Id}_P : P \rightarrow P$ and $b : P \rightarrow P$ are morphisms of categorical subspaces of B , so is their difference $\text{Id} - b : P \rightarrow P$. By Lemma 3.7, $S := [\text{Id} - b][P]$ is a categorical subspace of B whence of P . By Theorem 4.4, $I = (S) = ([\text{Id} - b][P])$ is a braided ideal of B . The conclusion follows by Lemma 4.2. $\square$

Lemma 4.6. *Let B be a braided bialgebra with infinitesimal part P . Let V be a categorical subspace of B generating B as an algebra and contained in P . Let $j : V \rightarrow P$ be the canonical inclusion. The following are equivalent for a K -linear map $b : P \rightarrow V$:*

- (i) $jb : P \rightarrow P$ is a morphism of categorical subspaces of B ;
- (ii) $b : P \rightarrow V$ fulfills

$$(4.1) \quad c_V(b \otimes V) = (V \otimes b)c_{P,V} \quad \text{and} \quad c_V(V \otimes b) = (b \otimes V)c_{V,P}.$$

Proof. (i) $\Rightarrow$ (ii) Clearly b fulfills (4.1) if jb fulfills (3.3).

(ii) $\Rightarrow$ (i) Since B is generated by V , we have that $B = \cup V^n$ where $V^n := V^{\cdot B^n}$ for all $n \in \mathbb{N}$. Hence, in order to prove that jb fulfills (3.3) it suffices to check, by induction on $n \geq 1$, that b fulfills

$$c_{V,V^n}(b \otimes V^n) = (V^n \otimes b) c_{P,V^n} \quad c_{V^n,V}(V^n \otimes b) = (b \otimes V^n) c_{V^n,P}.$$

Note that the notations above make sense in view of Lemma 3.5 and Lemma 3.9. For $n = 1$ there is nothing to prove as $c_{V,V} = c_V$. Assume that the formulas are true for $n - 1$. Then, we have

$$\begin{aligned} & c_{V,V^n}(b \otimes V^n) (P \otimes m_{V^{n-1},V}) \\ &= c_{V,V^n}(V \otimes m_{V^{n-1},V}) (b \otimes V^{n-1} \otimes V) \\ &\stackrel{(2.2)}{=} (m_{V^{n-1},V} \otimes V) (V^{n-1} \otimes c_V) (c_{V,V^{n-1}} \otimes V) (b \otimes V^{n-1} \otimes V) \\ &= (m_{V^{n-1},V} \otimes V) (V^{n-1} \otimes c_V) (V^{n-1} \otimes b \otimes V) (c_{P,V^{n-1}} \otimes V) \\ &= (m_{V^{n-1},V} \otimes V) (V^{n-1} \otimes V \otimes b) (V^{n-1} \otimes c_{P,V}) (c_{P,V^{n-1}} \otimes V) \\ &= (V^n \otimes b) (m_{V^{n-1},V} \otimes P) (V^{n-1} \otimes c_{P,V}) (c_{P,V^{n-1}} \otimes V) \\ &\stackrel{(2.2)}{=} (V^n \otimes b) c_{P,V^n} (P \otimes m_{V^{n-1},V}) \end{aligned}$$

where $m_{V^{n-1},V} : V^{n-1} \otimes V \rightarrow V^n$ is induced by the multiplication of B . Since $P \otimes m_{V^{n-1},V}$ is surjective, we arrive at $c_{V,V^n}(b \otimes V^n) = (V^n \otimes b) c_{P,V^n}$. Similarly one gets $c_{V^n,V}(V^n \otimes b) = (b \otimes V^n) c_{V^n,P}$. $\square$

Corollary 4.7. *Let B be a braided bialgebra with infinitesimal part P . Let V be a braided subspace of B generating B as an algebra and contained in P . Let $b : P \rightarrow V$ fulfill (4.1). Then*

$$W(B, b) := \frac{B}{([\text{Id} - b][P])}$$

carries a unique connected braided bialgebra structure such that the canonical projection $\pi_W : B \rightarrow W(B, b)$ is a braided bialgebra homomorphism.

Proof. It follows by Proposition 3.10, Lemma 4.6 and Proposition 4.5. $\square$

Definition 4.8. Let (V, c) be a braided vector space. A (V, c) -**bialgebra** is a pair (B, γ) such that

- B is a braided bialgebra with infinitesimal part (P, c_P) ;
- $\gamma : (V, c) \rightarrow (P, c_P)$ is a morphism of braided vector spaces;
- $\gamma(V)$ generates B as a K -algebra.

A **morphism** $\pi : (B, \gamma) \rightarrow (B', \gamma')$ **of (V, c) -bialgebras** is a braided bialgebra homomorphism $\pi : B \rightarrow B'$ such that $P(\pi)\gamma = \gamma'$ where $P(\pi) : P(B) \rightarrow P(B')$ is the obvious induced morphism.

Remark 4.9. Let (V, c) be a braided vector space. Then any (V, c) -bialgebra (B, γ) is connected. In fact, $\gamma(V) \subseteq P(B)$ generates B as a K -algebra. Hence B is primitively generated. Thus Proposition 2.9 applies.

Lemma 4.10. *Let (V, c) be a braided vector space. The following assertions are equivalent for a braided bialgebra B .*

- (1) *There is a map γ such that (B, γ) is a (V, c) -bialgebra.*
- (2) *There is a surjective braided bialgebra map $\varphi : T(V, c) \rightarrow B$.*

Proof. Set $T := T(V, c)$.

(1) $\Rightarrow$ (2) By the universal property of the braided tensor algebra, there is a unique braided bialgebra map $\varphi : T \rightarrow B$. Since $\gamma(V)$ generates B as a K -algebra, then φ is surjective.

(2) $\Rightarrow$ (1) Since φ is a braided bialgebra map, it preserves primitive elements, so that one has $\varphi(V) \subseteq \varphi(P(T)) \subseteq P(\varphi(T)) \subseteq P(B)$. Thus we can set γ to be the restriction of φ to V . Since φ is compatible with braidings, one has that γ is a morphism of braided vector spaces. Since φ is surjective, then $\gamma(V)$ generates B as a K -algebra. In other words (B, γ) is a (V, c) -bialgebra. $\square$

Theorem 4.11. *Let (V, c) be a braided vector space and let (B, γ) be a (V, c) -bialgebra endowed with a map $b : P(B) \rightarrow V$ such that*

$$(4.2) \quad c_{\gamma(V)}(\gamma b \otimes \gamma(V)) = (\gamma(V) \otimes \gamma b) c_{P(B), \gamma(V)},$$

$$(4.3) \quad c_{\gamma(V)}(\gamma(V) \otimes \gamma b) = (\gamma b \otimes \gamma(V)) c_{\gamma(V), P(B)}.$$

Then

$$W = W(B, \gamma b) = \frac{B}{([\text{Id} - \gamma b][P(B)])}$$

carries a unique (V, c) -bialgebra structure such that the canonical projection $\pi_W : B \rightarrow W$ is a (V, c) -bialgebra homomorphism.

Proof. $\gamma(V)$ is a braided subspace of B generating B as a K -algebra and contained in $P(B)$. In view of (4.2) and (4.3), we have that γb fulfills (4.1) where $\gamma(V)$ plays the role of V . By Corollary 4.7, $W(B, \gamma b)$ carries a unique braided bialgebra structure such that the canonical projection $\pi_W : B \rightarrow W$ is a braided bialgebra homomorphism. Clearly π_W induces a map $P(\pi_W) : P(B) \rightarrow P(W)$. Then $(W, P(\pi_W) \circ \gamma)$ is a (V, c) -bialgebra and, by construction, $\pi_W : B \rightarrow W$ is a (V, c) -bialgebra homomorphism. $\square$

Definition 4.12. With hypothesis and notations of Theorem 4.11, $(W, P(\pi_W) \circ \gamma)$ will be called the **quotient (V, c) -bialgebra of (B, γ) induced by b** .

5. UNIVERSAL ENVELOPING ALGEBRA

In this section we introduce and investigate the notion of braided Lie algebra for a braided vector space. We start by introducing the concept of bracket of finite rank and the corresponding universal enveloping algebra.

Definition 5.1. Let (V, c) be a braided vector space and let $n \in \mathbb{N}$.

Suppose there are (V, c) -bialgebras

$$(U^{[0]}, i^{[0]}), (U^{[1]}, i^{[1]}), \dots, (U^{[n]}, i^{[n]})$$

such that

- $U^{[0]} := T(V, c)$ and $i^{[0]} : V \rightarrow P(U^{[0]})$ is the restriction of the inclusion $V \rightarrow T(V, c)$;
- there exists a map $b^{[i]} : P(U^{[i]}) \rightarrow V$ such that (4.2) and (4.3) hold for $(B, \gamma, b) = (U^{[i]}, i^{[i]}, b^{[i]})$, for every $0 \leq i \leq n$;
- $(U^{[i]}, i^{[i]})$ is the quotient (V, c) -bialgebra of $(U^{[i-1]}, i^{[i-1]})$ induced by $b^{[i-1]}$ for every $1 \leq i \leq n$.

In this case we will say that

$$(b^{[i]})_{0 \leq i \leq n}$$

is a **bracket of rank n for a braided vector space (V, c)** .

Given a bracket $(b^{[i]})_{0 \leq i \leq n}$ of rank n for (V, c) we can define **the universal enveloping algebra of rank $n+1$** as the quotient (V, c) -bialgebra $(U^{[n+1]}, i^{[n+1]})$ of $(U^{[n]}, i^{[n]})$ induced by $b^{[n]}$ (see Definition 4.12). This will be denoted also by

$$U \left(V, c, (b^{[i]})_{0 \leq i \leq n} \right).$$

Note that, with this notation, we have that $U \left(V, c, (b^{[i]})_{0 \leq i \leq t-1} \right) = (U^{[t]}, i^{[t]})$ for all $1 \leq t \leq n$. This leads to define the **the universal enveloping algebra of rank 0** just as $U(V, c) := (U^{[0]}, i^{[0]})$.

For sake of simplicity we will adopt the following notation for all $n \in \mathbb{N}$

$$P^{[n]} := P(U^{[n]}), \quad V^{[n]} := i^{[n]}(V).$$

Hence for each $n \in \mathbb{N}$, (4.2) and (4.3) take form

$$(5.1) \quad c_{V^{[n]}} \left(i^{[n]} b^{[n]} \otimes V^{[n]} \right) = \left(V^{[n]} \otimes i^{[n]} b^{[n]} \right) c_{P^{[n]}, V^{[n]}},$$

$$(5.2) \quad c_{V^{[n]}} \left(V^{[n]} \otimes i^{[n]} b^{[n]} \right) = \left(i^{[n]} b^{[n]} \otimes V^{[n]} \right) c_{V^{[n]}, P^{[n]}},$$

and

$$U^{[n+1]} = \frac{U^{[n]}}{\left([\text{Id} - i^{[n]} b^{[n]}] [P^{[n]}] \right)}.$$

When there is matter of confusion, we will use the notations $P_V^{[n]} := P^{[n]}$, $U_V^{[n]} := U^{[n]}$, $b_V^{[n]} := b^{[n]}$, $i_V^{[n]} := i^{[n]}$ and $i_U^V := i_U$ as well.

We will denote by $\pi_n^{n+1} : U^{[n]} \rightarrow U^{[n+1]}$ the canonical projection for each $n \in \mathbb{N}$.

Next we give the definition of bracket for a braided vector space.

Definition 5.2. A **braided bracket** or simply a **bracket** for a braided vector space (V, c) is a family $b := (b^{[n]})_{n \in \mathbb{N}}$ of maps such that, for each $n \in \mathbb{N}$, the family $(b^{[i]})_{0 \leq i \leq n}$ is a bracket of rank n for (V, c) . This way we get a direct system of (V, c) -bialgebras

$$\left(U^{[0]}, i^{[0]} \right) \xrightarrow{\pi_0^1} \left(U^{[1]}, i^{[1]} \right) \xrightarrow{\pi_1^2} \left(U^{[2]}, i^{[2]} \right) \xrightarrow{\pi_2^3} \dots$$

The direct limit of the direct system of braided bialgebras

$$U^{[0]} \xrightarrow{\pi_0^1} U^{[1]} \xrightarrow{\pi_1^2} U^{[2]} \xrightarrow{\pi_2^3} \dots$$

will be denoted by

$$(U(V, c, b), \pi_U) := \left(U^{[\infty]}, \pi_n^\infty \right) := \varinjlim U \left(V, c, (b^{[i]})_{0 \leq i \leq n-1} \right).$$

By Lemma 2.12, $U(V, c, b)$ carries a natural braided bialgebra structure that makes it the direct limit of $\left((U^{[u]})_{u \in \mathbb{N}}, (\pi_u^v)_{u, v \in \mathbb{N}} \right)$ as a direct system of braided bialgebras. $U(V, c, b)$ will be called **the universal enveloping algebra of (V, c, b)** .

Denote by

$$i_U : V \rightarrow U(V, c, b)$$

the (not necessarily injective) canonical map and set $P^{[\infty]} := P(U^{[\infty]})$. Note that $\text{Im}(i_U) \subseteq P^{[\infty]}$ so that i_U induces a morphism of braided vector spaces $i^{[\infty]} : V \rightarrow P^{[\infty]}$.

Clearly $(U^{[\infty]}, i^{[\infty]})$ is a (V, c) -bialgebra.

When there is matter of confusion, we will use the notation $i_U^V := i_U$ as well.

Next aim is to describe the image of the map $i_U : V \rightarrow U(V, c, b)$.

Theorem 5.3. *Let (V, c) be a braided vector space endowed with a bracket b . Then $P^{[\infty]} = \varinjlim P^{[n]}$. Moreover $i^{[\infty]} : V \rightarrow P^{[\infty]}$ is surjective i.e.*

$$\text{Im}(i_U) = P(U(V, c, b)).$$

Proof. The braided bialgebra homomorphism $\pi_{n-1}^n := U^{[n-1]} \rightarrow U^{[n]}$ induces a direct system

$$P^{[0]} \xrightarrow{P(\pi_0^1)} P^{[1]} \xrightarrow{P(\pi_1^2)} P^{[2]} \xrightarrow{P(\pi_2^3)} \dots$$

Let us prove that $P^{[\infty]} = \varinjlim P^{[n]}$. Let $\delta^{[n]} := \delta_{U^{[n]}} : U^{[n]} \rightarrow U^{[n]} \otimes U^{[n]}$ be defined as in 2.5. Then

$$\begin{aligned} (\pi_{n-1}^n \otimes \pi_{n-1}^n) \delta^{[n-1]}(x) &= (\pi_{n-1}^n \otimes \pi_{n-1}^n) (x \otimes 1_{U^{[n-1]}} + 1_{U^{[n-1]}} \otimes x - \Delta_{U^{[n-1]}}(x)) \\ &= \pi_{n-1}^n(x) \otimes 1_{U^{[n]}} + 1_{U^{[n]}} \otimes \pi_{n-1}^n(x) - (\pi_{n-1}^n \otimes \pi_{n-1}^n) \Delta_{U^{[n-1]}}(x) \\ &= \pi_{n-1}^n(x) \otimes 1_{U^{[n]}} + 1_{U^{[n]}} \otimes \pi_{n-1}^n(x) - \Delta_{U^{[n]}} \pi_{n-1}^n(x) \\ &= \delta^{[n]} \pi_{n-1}^n(x) \end{aligned}$$

so that $(\pi_{n-1}^n \otimes \pi_{n-1}^n) \delta^{[n-1]} = \delta^{[n]} \pi_{n-1}^n$. Hence from the exact sequence

$$0 \rightarrow P^{[n]} \rightarrow U^{[n]} \xrightarrow{\delta^{[n]}} U^{[n]} \otimes U^{[n]}$$

we get the exact sequence

$$0 \rightarrow \varinjlim P^{[n]} \rightarrow \varinjlim U^{[n]} \xrightarrow{\varinjlim \delta^{[n]}} \varinjlim (U^{[n]} \otimes U^{[n]})$$

which can be rewritten as

$$0 \rightarrow \varinjlim P^{[n]} \rightarrow U^{[\infty]} \xrightarrow{\delta^{[\infty]}} U^{[\infty]} \otimes U^{[\infty]}$$

so that $P^{[\infty]} = \varinjlim P^{[n]}$. For each $z \in P^{[\infty]}$, there is $t \in \mathbb{N}$ and $z_t \in P^{[t]}$ such that $z = \pi_t^\infty(z_t)$. Thus

$$z = \pi_t^\infty(z_t) = \pi_{t+1}^\infty \pi_t^{t+1}(z_t) \stackrel{z_t \in P^{[t]}}{=} \pi_{t+1}^\infty \pi_t^{t+1} i^{[t]} b^{[t]}(z_t) = i^{[\infty]} b^{[t]}(z_t) \in \text{Im}(i^{[\infty]})$$

whence $P^{[\infty]} \subseteq \text{Im}(i^{[\infty]})$. Since $\text{Im}(i^{[\infty]}) \subseteq P^{[\infty]}$ we conclude. $\square$

The map $i_U : V \rightarrow U(V, c, b)$ needs not to be injective in general. Thus it makes sense to introduce the following definition.

Definition 5.4. We say that (V, c, b) is a **braided Lie algebra** whenever

- (V, c) is a braided vector space;
- b is a bracket on (V, c) ;
- $i_U : V \rightarrow U(V, c, b)$ is injective (**implicit Jacobi identity**).

Remark 5.5. The third condition above, was called implicit Jacobi identity having in mind [AMS, Theorem 4.3].

The following result will be crucial in proving the main theorem.

Corollary 5.6. *Let (V, c, b) be a braided Lie algebra. The canonical map $i_U : V \rightarrow U(V, c, b)$ induces an isomorphism of braided vector spaces between V and $P(U(V, c, b))$ namely the map $i^{[\infty]} : V \rightarrow P^{[\infty]}$.*

Proof. Injectivity of $i_U : V \rightarrow U(V, c, b)$ implies Injectivity of $i^{[\infty]} : V \rightarrow P^{[\infty]}$. The conclusion follows by Theorem 5.3. $\square$

We now give a different characterization of the implicit Jacobi identity.

Proposition 5.7. *Let (V, c) is a braided vector space endowed with a bracket b . The following assertions are equivalent.*

- (i) *the implicit Jacobi identity holds i.e. (V, c, b) is a braided Lie algebra.*
- (ii) *$b^{[n]} i^{[n]} = \text{Id}_V$ for all $n \in \mathbb{N}$.*
- (iii) *$b^{[0]} i^{[0]} = \text{Id}_V$ and $b^{[n+1]} P(\pi_n^{n+1}) = b^{[n]}$ for all $n \in \mathbb{N}$.*

Here $P(\pi_n^{n+1}) : P^{[n]} \rightarrow P^{[n+1]}$ denotes the map induced by π_n^{n+1} .

Proof. (i) $\Rightarrow$ (ii) We have

$$\begin{aligned} & i_U (\text{Id}_V - b^{[n]} i^{[n]}) (V) \\ &= P(\pi_n^\infty) i^{[n]} (\text{Id}_V - b^{[n]} i^{[n]}) (V) \\ &= \pi_n^\infty (\text{Id}_{U^{[n]}} - i^{[n]} b^{[n]}) i^{[n]} (V) \subseteq \pi_{n+1}^\infty \pi_n^{n+1} (\text{Id}_{P^{[n]}} - i^{[n]} b^{[n]}) [P^{[n]}] = 0 \end{aligned}$$

so that $i_U (\text{Id}_V - b^{[n]} i^{[n]}) = 0$. Then injectivity of i_U yields $b^{[n]} i^{[n]} = \text{Id}_V$.

(ii) $\Rightarrow$ (i) Since $i^{[n]} : V \rightarrow P^{[n]}$ is injective for all $n \in \mathbb{N}$, we get that $i^{[\infty]} = \varinjlim i^{[n]} : V \rightarrow \varinjlim P^{[n]} = P^{[\infty]}$ is injective too. Since $i^{[\infty]}$ is obtained by corestricting i_U to $P^{[\infty]}$, we have that i_U is injective too.

(i) $\Rightarrow$ (iii) For all $x \in P^{[n]}$ we have

$$\begin{aligned}
 i_U \left[b^{[n]} - b^{[n+1]} P(\pi_n^{n+1}) \right] (x) &= \left[i_U b^{[n]} - i_U b^{[n+1]} P(\pi_n^{n+1}) \right] (x) \\
 &= \pi_{n+1}^\infty \pi_n^{n+1} i^{[n]} b^{[n]} (x) - \pi_{n+2}^\infty \pi_{n+1}^{n+2} i^{[n+1]} b^{[n+1]} P(\pi_n^{n+1}) (x) \\
 &= \pi_{n+1}^\infty \pi_n^{n+1} (x) - \pi_{n+2}^\infty \pi_{n+1}^{n+2} P(\pi_n^{n+1}) (x) \\
 &= \pi_n^\infty (x) - \pi_{n+1}^\infty P(\pi_n^{n+1}) (x) \\
 &= \pi_n^\infty (x) - \pi_{n+1}^\infty \pi_n^{n+1} (x) = 0
 \end{aligned}$$

so that $i_U [b^{[n]} - b^{[n+1]} P(\pi_n^{n+1})] = 0$. Injectivity of i_U implies $b^{[n]} = b^{[n+1]} P(\pi_n^{n+1})$. By (i) $\Rightarrow$ (ii) we have that $b^{[0]} i^{[0]} = \text{Id}_V$.

(iii) $\Rightarrow$ (ii) We prove it by induction on $n \in \mathbb{N}$. For $n = 0$ there is nothing to prove. By definition of $i^{[n+1]}$ and by induction hypothesis we get

$$b^{[n+1]} i^{[n+1]} = b^{[n+1]} P(\pi_n^{n+1}) i^{[n]} = b^{[n]} i^{[n]} = \text{Id}_V.$$

□

In view of the previous result, we are able to give an alternative description of the universal enveloping algebra of positive rank.

Corollary 5.8. *Let (V, c, b) be a braided Lie algebra. Then*

$$U^{[n+1]} = \frac{U^{[n]}}{(\text{Ker}(b^{[n]}))}, \text{ for each } n \in \mathbb{N}.$$

Proof. By Proposition 5.7, we have that $b_V^{[n]} i_V^{[n]} = \text{Id}_V$ for each $n \in \mathbb{N}$. This condition entails that $\text{Im}(\text{Id} - i_V^{[n]} b_V^{[n]}) = \text{Ker}(b^{[n]})$. □

Next aim is to show that the universal enveloping algebra introduced in [Ar2] can be understood as a universal enveloping algebra of rank one in the sense of the present paper.

Remark 5.9. Let (V, c) be a braided vector space. Denote by $E(V, c)$ the space spanned by primitive elements of $T(V, c)$ of homogeneous degree at least two. In [Ar2, Definition 3.2], a bracket for (V, c) was defined as a map $\beta : E(V, c) \rightarrow V$ such that

$$c(\beta \otimes V) = (V \otimes \beta) c_{E(V, c), V}, \quad c(V \otimes \beta) = (\beta \otimes V) c_{V, E(V, c)}.$$

The corresponding universal enveloping algebra was defined in [Ar2, Definition 3.5] as

$$\mathbb{U}(V, c, \beta) = \frac{T(V, c)}{((\text{Id} - \beta)[E(V, c)])}.$$

When the canonical map $i_{\mathbb{U}} : V \rightarrow \mathbb{U}(V, c, \beta)$ is injective, the tern (V, c, β) was called a braided Lie algebra in [Ar2, Definition 4.1]. Now the primitive part of $\mathbb{U}(V, c, \beta)$ identifies with (V, c) itself if (V, c) is in the class $\mathcal{S}$ consisting of those braided vector spaces such that the kernel of the canonical projection $p_0^\infty : T(V, c) \rightarrow \mathcal{B}(V, c)$, onto the Nichols algebra $\mathcal{B}(V, c)$, is the two-sided ideal of $T(V, c)$ generated by $E(V, c)$, see [Ar2, Corollary 5.5]. This property implies that all primitively generated braided bialgebras whose infinitesimal part is in $\mathcal{S}$ are universal enveloping algebras in the above sense of their primitive part, see [Ar2, Theorem 5.7]. Significant braided vector spaces belong to the class $\mathcal{S}$, see [Ar1], where elements in $\mathcal{S}$ are called braided vector spaces of strongness degree at most one. Anyway, the fact that not all braided vector spaces are in $\mathcal{S}$, led us to introduce in the present paper a different notion of braided Lie algebra and universal enveloping algebra. Next result shows that the old definition can be understood as a first step of the new one.

Theorem 5.10. *Let (V, c, b) be a braided Lie algebra. Let $\beta : E(V, c) \rightarrow V$ be the restriction of $b^{[0]}$ to $E(V, c)$. Then (V, c, β) is a braided Lie algebra in the sense of Remark 5.9 and $\mathbb{U}(V, c, \beta) = U(V, c, b^{[0]})$. Moreover, if the braided vector space (V, c) is in the class $\mathcal{S}$, then $U(V, c, b) = U(V, c, b^{[0]})$.*

Proof. Since $i^{[0]} : V \rightarrow P^{[0]}$ is injective, V identifies with $V^{[0]} = i^{[0]}(V)$ so that (5.1) and (5.2) in case $n = 0$ rewrites as

$$c(b^{[0]} \otimes V) = (V \otimes b^{[0]}) c_{P^{[0]}, V}, \quad c(V \otimes b^{[0]}) = (b^{[0]} \otimes V) c_{V, P^{[0]}}.$$

Restricting the former equality to $E(V, c) \otimes V$ and the latter to $V \otimes E(V, c)$ yields

$$c(\beta \otimes V) = (V \otimes \beta) c_{E(V, c), V}, \quad c(V \otimes \beta) = (\beta \otimes V) c_{V, E(V, c)}$$

which means that β is a bracket in the sense of Remark 5.9. By [Ar2, Remark 2.9], $P^{[0]} = P(U^{[0]}) = V^{[0]} \oplus E(V, c)$. By Proposition 5.7, $b^{[n]} \circ i^{[n]} = \text{Id}_V$ for all $n \in \mathbb{N}$ so that we have

$$\begin{aligned} (\text{Id} - i^{[0]}b^{[0]}) [P^{[0]}] &= (\text{Id} - i^{[0]}b^{[0]}) [V^{[0]} \oplus E(V, c)] \\ &= (\text{Id} - i^{[0]}b^{[0]}) [V^{[0]}] + (\text{Id} - i^{[0]}b^{[0]}) [E(V, c)] \\ &= (\text{Id} - i^{[0]}b^{[0]}) i^{[0]}(V) + (\text{Id} - i^{[0]}b^{[0]}) [E(V, c)] \\ &= (\text{Id} - \beta) [E(V, c)] \end{aligned}$$

whence

$$U(V, c, b^{[0]}) = \frac{U^{[0]}}{((\text{Id} - i^{[0]}b^{[0]}) [P^{[0]}])} = \frac{T(V, c)}{((\text{Id} - \beta) [E(V, c)])} = \mathbb{U}(V, c, \beta).$$

Now, the canonical map $i_{\mathbb{U}} : V \rightarrow \mathbb{U}(V, c, \beta)$ corestricts to the map $i^{[1]} : V \rightarrow P^{[1]}$ which is split injective by $b^{[1]}$. Thus $i_{\mathbb{U}}$ is injective that is (V, c, β) is a braided Lie algebra in the sense of Remark 5.9.

Assume now the braided vector space (V, c) is in $\mathcal{S}$. Then the space of primitive elements in $\mathbb{U}(V, c, \beta)$ identifies with V via $i_{\mathbb{U}}$. Therefore, the projection $\pi_1^\infty : U(V, c, b^{[0]}) \rightarrow U(V, c, b)$ is injective on primitive elements of $U(V, c, b^{[0]})$. By [Mo, Lemma 5.3.3] π_1^∞ is injective whence bijective. $\square$

Lemma 5.11. *Let (V, c, b) be a braided Lie algebra such that $U(V, c, b) = U(V, c, (b^{[t]})_{0 \leq t \leq n})$ for some $n \in \mathbb{N}$. Then, for all $t \geq n + 1$, $b^{[t]} = b^{[n+1]}$. Moreover $b^{[n+1]}$ is bijective and it is the inverse of $i^{[n+1]} : V \rightarrow P^{[n+1]}$.*

Proof. Let $t \geq n + 1$. By assumption, the canonical map $\pi_{n+1}^\infty : U^{[n+1]} \rightarrow U^{[\infty]}$ is the identity so that $\pi_t^\infty : U^{[t]} \rightarrow U^{[\infty]}$ is the identity too. Now, by Corollary 5.6, we know that $i^{[\infty]}$ is bijective. By construction, $i^{[\infty]} = P(\pi_t^\infty) i^{[t]} = i^{[t]}$. In particular $i^{[t]}$ is bijective and $i^{[t]} = i^{[n+1]}$. From Proposition 5.7, we know that $b^{[t]} i^{[t]} = \text{Id}_V$. Thus $b^{[t]}$ is the inverse of $i^{[n+1]} : V \rightarrow P^{[n+1]}$ which is $b^{[n+1]}$. $\square$

6. THE MAIN RESULT

In this section we associates a braided Lie algebra structure to the infinitesimal part of a braided bialgebra. The braided bialgebra will then be recovered as the universal enveloping algebra of its infinitesimal braided Lie algebra.

Proposition 6.1. *Any braided algebra (A, c) admits a canonical bracket $b = (b^{[n]})_{n \in \mathbb{N}}$ such that (A, c, b) is a braided Lie algebra.*

Proof. Let us define recursively maps $b^{[n]} : P^{[n]} \rightarrow A$ such that $b^{[n]} i^{[n]} = \text{Id}_A$ and (5.1) and (5.2) hold for $V = A$.

$n = 0$) By the universal property of $U^{[0]} = T(A, c)$ (see [AMS, Theorem 1.17]), Id_A can be lifted to a unique braided algebra homomorphism $\varphi^{[0]} : U^{[0]} \rightarrow A$. Let $b^{[0]} : P^{[0]} \rightarrow A$ be the restriction of $\varphi^{[0]}$ to $P^{[0]}$. Clearly $b^{[0]} \circ i^{[0]} = \text{Id}_A$. Let us check that

$$(6.1) \quad c_{U^{[0]}, A^{[0]}} (i^{[0]} \varphi^{[0]} \otimes A^{[0]}) = (A^{[0]} \otimes i^{[0]} \varphi^{[0]}) c_{U^{[0]}, A^{[0]}}$$

holds (here $A^{[0]} := i^{[0]}(A)$). Note that the notation $c_{U^{[0]}, A^{[0]}}$ makes sense as, by Proposition 3.10, $A^{[0]}$ is a categorical subspace of $(U^{[0]}, c_{U^{[0]}})$. Let $j_t : A^{\otimes t} \rightarrow U^{[0]}$ be the canonical injection. Then

$$\begin{aligned}
& c_{U^{[0]}, A^{[0]}} \left(i^{[0]} \varphi^{[0]} \otimes A^{[0]} \right) (j_t \otimes j_0) \\
&= c_{U^{[0]}, A^{[0]}} \left(i^{[0]} \varphi^{[0]} j_t \otimes j_0 \right) \\
&= c_{U^{[0]}, A^{[0]}} \left(i^{[0]} \varphi^{[0]} j_t \otimes i^{[0]} \right) \\
&= \left(i^{[0]} \otimes i^{[0]} \right) c_A \left(\varphi^{[0]} j_t \otimes A \right) \\
&= \left(i^{[0]} \otimes i^{[0]} \right) c_A \left(m_A^{t-1} \otimes A \right) \\
&\stackrel{(2.2)}{=} \left(i^{[0]} \otimes i^{[0]} \right) \left(A \otimes m_A^{t-1} \right) c_{A^{\otimes t}, A} \\
&= \left(i^{[0]} \otimes i^{[0]} \right) \left(A \otimes \varphi^{[0]} j_t \right) c_{A^{\otimes t}, A} \\
&= \left(A^{[0]} \otimes i^{[0]} \varphi^{[0]} \right) (j_0 \otimes j_t) c_{A^{\otimes t}, A} \\
&= \left(A^{[0]} \otimes i^{[0]} \varphi^{[0]} \right) c_{U^{[0]}, U^{[0]}} (j_t \otimes j_0) \\
&= \left(A^{[0]} \otimes i^{[0]} \varphi^{[0]} \right) c_{U^{[0]}, A^{[0]}} (j_t \otimes j_0).
\end{aligned}$$

Since this equality holds for all $t \in \mathbb{N}$, and $A^{[0]} = \text{Im}(i^{[0]}) = \text{Im}(j_0)$, we get (6.1). Now we can prove that (5.1) is fulfilled for $n = 0$. In fact, for each $x \in P^{[0]}$ and $a \in A^{[0]}$ we have

$$\begin{aligned}
& c_{A^{[0]}} \left(i^{[0]} b^{[0]} \otimes A^{[0]} \right) (x \otimes a) \\
&= c_{U^{[0]}, A^{[0]}} \left(i^{[0]} \varphi^{[0]} \otimes A^{[0]} \right) (x \otimes a) \\
&\stackrel{(6.1)}{=} \left(A^{[0]} \otimes i^{[0]} \varphi^{[0]} \right) c_{U^{[0]}, A^{[0]}} (x \otimes a) \\
&= \left(A^{[0]} \otimes i^{[0]} b^{[0]} \right) c_{P^{[0]}, A^{[0]}} (x \otimes a).
\end{aligned}$$

Similarly one proves (5.2).

$n \geq 1$) We have

$$\begin{aligned}
& \varphi^{[n-1]} \left(\text{Id} - i^{[n-1]} b^{[n-1]} \right) \left[P^{[n-1]} \right] \\
&= \left(\varphi^{[n-1]} - \varphi^{[n-1]} i^{[n-1]} b^{[n-1]} \right) \left[P^{[n-1]} \right] \\
&= \left(\varphi^{[n-1]} - b^{[n-1]} \right) \left[P^{[n-1]} \right] = \left(\varphi^{[n-1]} - \varphi^{[n-1]} \right) \left[P^{[n-1]} \right] = 0
\end{aligned}$$

so that $\varphi^{[n-1]} : U^{[n-1]} \rightarrow A$ quotients to a map $\varphi^{[n]} : U^{[n]} \rightarrow A$. Let $b^{[n]} : P^{[n]} \rightarrow A$ be the restriction of $\varphi^{[n]}$ to $P^{[n]}$. Clearly $b^{[n]} i^{[n]} = \text{Id}_A$. Let us prove that (5.1) holds. Let $x \in P^{[n]}$ and $y \in A^{[n]}$. Then there is $x_A \in U^{[0]}$ such that $x = \pi_0^n(x_A)$ and $y_A \in A$ such that $y = i^{[n]}(y_A) = \pi_0^n i^{[0]}(y_A)$. Whence

$$\begin{aligned}
& c_{A^{[n]}} \left(i^{[n]} b^{[n]} \otimes A^{[n]} \right) (x \otimes y) \\
&= c_{U^{[n]}} \left(i^{[n]} b^{[n]} \otimes A^{[n]} \right) \left(\pi_0^n(x_A) \otimes \pi_0^n i^{[0]}(y_A) \right) \\
&= c_{U^{[n]}} \left(i^{[n]} \varphi^{[n]} \pi_0^n(x_A) \otimes \pi_0^n i^{[0]}(y_A) \right) \\
&= c_{U^{[n]}} \left(\pi_0^n i^{[0]} \varphi^{[0]}(x_A) \otimes \pi_0^n i^{[0]}(y_A) \right) \\
&= (\pi_0^n \otimes \pi_0^n) c_{U^{[0]}} \left(i^{[0]} \varphi^{[0]} \otimes U^{[0]} \right) (x_A \otimes i^{[0]}(y_A))
\end{aligned}$$

$$\begin{aligned}
&= (\pi_0^n \otimes \pi_0^n) c_{U^{[0]}, A^{[0]}} \left(i^{[0]} \varphi^{[0]} \otimes A^{[0]} \right) \left(x_A \otimes i^{[0]}(y_A) \right) \\
&\stackrel{(6.1)}{=} (\pi_0^n \otimes \pi_0^n) \left(A^{[0]} \otimes i^{[0]} \varphi^{[0]} \right) c_{U^{[0]}, A^{[0]}} \left(x_A \otimes i^{[0]}(y_A) \right) \\
&= \left(\pi_0^n \otimes \pi_0^n i^{[0]} \varphi^{[0]} \right) c_{U^{[0]}, A^{[0]}} \left(x_A \otimes i^{[0]}(y_A) \right) \\
&= \left(\pi_0^n \otimes i^{[n]} \varphi^{[0]} \right) c_{U^{[0]}} \left(x_A \otimes i^{[0]}(y_A) \right) \\
&= \left(\pi_0^n \otimes i^{[n]} \varphi^{[n]} \pi_0^n \right) c_{U^{[0]}} \left(x_A \otimes i^{[0]}(y_A) \right) \\
&= \left(A^{[n]} \otimes i^{[n]} b^{[n]} \right) c_{U^{[n]}} \left(\pi_0^n(x_A) \otimes \pi_0^n i^{[0]}(y_A) \right) \\
&= \left(A^{[n]} \otimes i^{[n]} b^{[n]} \right) c_{P^{[n]}, A^{[n]}} (x \otimes y)
\end{aligned}$$

Similarly one proves (5.2). We have so proved that $b := (b^{[n]})_{n \in \mathbb{N}}$ is a bracket for (A, c) . Since, $b^{[n]} i^{[n]} = \text{Id}_A$ for each $n \in \mathbb{N}$, by Proposition 5.7, we conclude that (A, c, b) is a braided Lie algebra. $\square$

6.2. Let $h : (V, c_V) \rightarrow (W, c_W)$ be a morphism of braided vector spaces. Assume that (V, c_V) and (W, c_W) are endowed with a bracket b_V and b_W respectively. By the universal property of the tensor algebra, h induces a braided bialgebra homomorphism

$$U_h^{[0]} : U_V^{[0]} \rightarrow U_W^{[0]}$$

such that $U_h^{[0]} i_V^{[0]} = i_W^{[0]} h$ where $i_W^{[0]}$ is thought as a map into $U_W^{[0]}$ and $i_V^{[0]}$ as a map into $U_V^{[0]}$. The map $U_h^{[0]}$ preserves primitives so that it gives rise to a map

$$P_h^{[0]} : P_V^{[0]} \rightarrow P_W^{[0]}$$

such that $P_h^{[0]} i_V^{[0]} = i_W^{[0]} h$. Let $\pi_0^1 : U_V^{[0]} \rightarrow U_V^{[1]}$ and $\Pi_0^1 : U_W^{[0]} \rightarrow U_W^{[1]}$ be the canonical projections. Suppose

$$(6.2) \quad h b_V^{[n]} = b_W^{[n]} P_h^{[n]}$$

holds for $n = 0$. Since

$$\begin{aligned}
\Pi_0^1 U_h^{[0]} \left(\text{Id} - i_V^{[0]} b_V^{[0]} \right) \left[P_V^{[0]} \right] &= \Pi_0^1 \left(U_h^{[0]} - U_h^{[0]} i_V^{[0]} b_V^{[0]} \right) \left[P_V^{[0]} \right] \\
&= \Pi_0^1 \left(P_h^{[0]} - i_W^{[0]} h b_V^{[0]} \right) \left[P_V^{[0]} \right] \\
&= \Pi_0^1 \left(P_h^{[0]} - i_W^{[0]} b_W^{[0]} P_h^{[0]} \right) \left[P_V^{[0]} \right] \\
&= \Pi_0^1 \left(\text{Id} - i_W^{[0]} b_W^{[0]} \right) P_h^{[0]} \left[P_V^{[0]} \right] \subseteq \Pi_0^1 \left(\text{Id} - i_W^{[0]} b_W^{[0]} \right) \left[P_W^{[0]} \right] = 0
\end{aligned}$$

we have that $U_h^{[0]}$ induces a map $U_h^{[1]} : U_V^{[1]} \rightarrow U_W^{[1]}$ such that $U_h^{[1]} \pi_0^1 = \Pi_0^1 U_h^{[0]}$. Using that Π_0^1 is a surjective braided bialgebra homomorphism, one can check that $U_h^{[1]}$ is a braided bialgebra homomorphism too. Note that

$$U_h^{[1]} i_V^{[1]} = U_h^{[1]} \pi_0^1 i_V^{[0]} = \Pi_0^1 U_h^{[0]} i_V^{[0]} = \Pi_0^1 i_W^{[0]} h = i_W^{[1]} h.$$

As in the case of $U_h^{[0]}$, the map $U_h^{[1]}$ preserves primitives so that it gives rise to a map

$$P_h^{[1]} : P_V^{[1]} \rightarrow P_W^{[1]}.$$

Now supposing (6.2) holds for $n = 1$, one can define $P_h^{[2]}$ and so on.

Definition 6.3. A morphism of braided brackets or simply a morphism of brackets $h : (V, c_V, b_V) \rightarrow (W, c_W, b_W)$ is a morphism of braided vector spaces $h : (V, c_V) \rightarrow (W, c_W)$ such that (6.2) holds for every $n \in \mathbb{N}$, where $P_h^{[n]}$ is constructed as in 6.2. Note that $P_h^{[0]}$ always exists while, for $t \geq 1$, $P_h^{[t]}$ exists as (6.2) holds for $n = t - 1$.

Lemma 6.4. *Let (W, c_W, b_W) a braided Lie algebra and let $h : (V, c_V) \rightarrow (W, c_W)$ be an injective morphism of braided vector spaces. If, for each $n \in \mathbb{N}$, there exists a K -linear map $b_V^{[n]} : P_V^{[n]} \rightarrow V$ such that (6.2) holds, where $P_h^{[n]} : P_V^{[n]} \rightarrow P_W^{[n]}$ is constructed step by step as in 6.2, then (V, c_V, b_V) is a braided Lie algebra too and h is a morphism of braided brackets.*

Proof. Construct $U_h^{[0]} : U_V^{[0]} \rightarrow U_W^{[0]}$ and $P_h^{[0]} : P_V^{[0]} \rightarrow P_W^{[0]}$ as in 6.2. Note that $U_h^{[0]}$ restricts to a map $h^{[0]} : V^{[0]} \rightarrow W^{[0]}$ such that $h^{[0]}i_V^{[0]} = i_W^{[0]}h$ where $i_V^{[0]}$ is thought as a map into $V^{[0]}$ and $i_W^{[0]}$ as a map into $W^{[0]}$. For all $w \in P_V^{[0]}$ and $y \in V^{[0]}$ we have

$$\begin{aligned} & c_{P_W^{[0]}, W^{[0]}} \left(P_h^{[0]} \otimes h^{[0]} \right) (w \otimes y) \\ &= c_{U_W^{[0]}, U_W^{[0]}} \left(U_h^{[0]} \otimes U_h^{[0]} \right) (w \otimes y) \\ &= \left(U_h^{[0]} \otimes U_h^{[0]} \right) c_{U_V^{[0]}, U_V^{[0]}} (w \otimes y) = \left(h^{[0]} \otimes P_h^{[0]} \right) c_{P_V^{[0]}, V^{[0]}} (w \otimes y). \end{aligned}$$

Thus we have

$$(6.3) \quad c_{P_W^{[0]}, W^{[0]}} \left(P_h^{[0]} \otimes h^{[0]} \right) = \left(h^{[0]} \otimes P_h^{[0]} \right) c_{P_V^{[0]}, V^{[0]}}.$$

Hence, we have

$$\begin{aligned} & \left(h^{[0]} \otimes h^{[0]} \right) c_{V^{[n]}} \left(i_V^{[0]} b_V^{[0]} \otimes V^{[0]} \right) \\ &= c_{W^{[0]}} \left(h^{[0]} \otimes h^{[0]} \right) \left(i_V^{[0]} b_V^{[0]} \otimes V^{[0]} \right) \\ &= c_{W^{[0]}} \left(i_W^{[0]} h b_V^{[0]} \otimes h^{[0]} \right) \\ &\stackrel{(6.2)}{=} c_{W^{[0]}} \left(i_W^{[0]} b_W^{[0]} P_h^{[0]} \otimes h^{[0]} \right) \\ &= c_{W^{[0]}} \left(i_W^{[0]} b_W^{[0]} \otimes W^{[0]} \right) \left(P_h^{[0]} \otimes h^{[0]} \right) \\ &\stackrel{(5.1)}{=} \left(W^{[0]} \otimes i_W^{[0]} b_W^{[0]} \right) c_{P_W^{[0]}, W^{[0]}} \left(P_h^{[0]} \otimes h^{[0]} \right) \\ &\stackrel{(6.3)}{=} \left(h^{[0]} \otimes i_W^{[0]} b_W^{[0]} P_h^{[0]} \right) c_{P_V^{[0]}, V^{[0]}} \\ &\stackrel{(6.2)}{=} \left(h^{[0]} \otimes i_W^{[0]} h b_V^{[0]} \right) c_{P_V^{[0]}, V^{[0]}} \\ &= \left(h^{[0]} \otimes h^{[0]} \right) \left(V^{[0]} \otimes i_V^{[0]} b_V^{[0]} \right) c_{P_V^{[0]}, V^{[0]}} \end{aligned}$$

so that

$$\left(h^{[0]} \otimes h^{[0]} \right) c_{V^{[n]}} \left(i_V^{[0]} b_V^{[0]} \otimes V^{[0]} \right) = \left(h^{[0]} \otimes h^{[0]} \right) \left(V^{[0]} \otimes i_V^{[0]} b_V^{[0]} \right) c_{P_V^{[0]}, V^{[0]}}.$$

Now, since (W, c_W, b_W) is a braided Lie algebra, by Proposition 5.7, we have that $i_W^{[0]}$ is split injective. From this fact, injectivity of h and equality $h^{[0]}i_V^{[0]} = i_W^{[0]}h$, one easily gets that $\text{Ker}(h^{[0]}) = 0$ so that $h^{[0]}$ is injective. Thus, from the last displayed equality, we infer that $b_V^{[0]}$ fulfills (5.1). Similarly (5.2) is satisfied. Hence $b_V^{[0]}$ is a bracket for $U_V^{[0]}$ and one can construct $U_V^{[1]}$. Thus we can construct $U_h^{[1]} : U_V^{[1]} \rightarrow U_W^{[1]}$ and $P_h^{[1]} : P_V^{[1]} \rightarrow P_W^{[1]}$ as in 6.2 and $h^{[1]} : V^{[1]} \rightarrow W^{[1]}$ as we did for $h^{[0]}$ above. Then one can proceed as in the case of $U_h^{[0]}$ to prove that $b_V^{[1]}$ fulfills (5.1) and (5.2). Going on this way, we get that $b_V = \left(b_V^{[n]} \right)_{n \in \mathbb{N}}$ is a bracket on (V, c_V) . By (6.2), h is a morphism of braided brackets. Now, let

$$U_h^{[\infty]} := \varinjlim U_h^{[n]} : U(V, c_V, b_V) = U_V^{[\infty]} \rightarrow U_W^{[\infty]} = U(W, c_W, b_W).$$

Denoting by $i_U^V : V \rightarrow U(V, c_V, b_V)$ and $i_U^W : W \rightarrow U(W, c_W, b_W)$ the canonical maps, we get that $U_h^{[\infty]}i_U^V = i_U^W h$. Since both i_U^W and h are injective, we deduce that i_U^V is injective too. Therefore (V, c_V, b_V) is a braided Lie algebra. $\square$

Theorem 6.5. *Let A be a braided bialgebra with infinitesimal part (P, c_P) . Then the braided Lie algebra structure of A of Proposition 6.1 induces a braided Lie algebra structure (P, c_P, b_P) .*

Proof. Let $\varphi^{[n]} : U_A^{[n]} \rightarrow A$ be defined as in the proof of Proposition 6.1. In order to avoid ambiguities on the notation $P^{[n]}$, we set $V := P$. By Proposition 6.1, A has a braided Lie algebra structure, say (A, c_A, b_A) , and by Proposition 5.7, $b_A^{[n]} i_A^{[n]} = \text{Id}_A$ for all $n \in \mathbb{N}$. The canonical injection $\lambda : V \rightarrow A$ is a morphism of braided vector spaces. We will apply Lemma 6.4 and use the notations in its proof. For any $n \in \mathbb{N}$, consider the map

$$b_A^{[n]} P_\lambda^{[n]} : P_V^{[n]} \rightarrow A.$$

By construction of $b_A^{[n]}$, for $x \in P_V^{[n]}$ we have

$$b_A^{[n]} P_\lambda^{[n]}(x) = \varphi^{[n]} P_\lambda^{[n]}(x) = \varphi^{[n]} U_\lambda^{[n]}(x).$$

Since $\varphi^{[n]} U_\lambda^{[n]}$ is a braided bialgebra homomorphism, it preserves primitive elements so that $\varphi^{[n]} U_\lambda^{[n]}(x) \in V$. Thus $b_A^{[n]} P_\lambda^{[n]}(x) \in V$ whence there exists a unique map $b_V^{[n]} : P_V^{[n]} \rightarrow V$ such that $\lambda b_V^{[n]} = b_A^{[n]} P_\lambda^{[n]}$. By Lemma 6.4, (V, c_V, b_V) is a braided Lie algebra too. $\square$

Definition 6.6. With the same assumptions and notations as in Theorem 6.5, (P, c_P, b_P) will be called the **infinitesimal braided Lie algebra of A** .

We are now able to state the universal property of the universal enveloping algebra.

Theorem 6.7 (The universal property). *Let A be a braided bialgebra with infinitesimal braided Lie algebra (P, c_P, b_P) . Then every morphism of braided brackets $h : (V, c, b) \rightarrow (P, c_P, b_P)$ can be lifted to a morphism of braided bialgebras $\bar{h} : U(V, c, b) \rightarrow A$.*

Proof. Let $\varphi^{[n]} : U_A^{[n]} \rightarrow A$ be defined as in the proof of Proposition 6.1. Let $\lambda : P \rightarrow A$ be the canonical map. Set $h^{[n]} := \varphi^{[n]} U_{\lambda h}^{[n]} : U_V^{[n]} \rightarrow A$. Let $\pi_{n-1}^{[n]} : U_V^{[n-1]} \rightarrow U_V^{[n]}$ and $\Pi_{n-1}^{[n]} : U_A^{[n-1]} \rightarrow U_A^{[n]}$ be the canonical projections. We have

$$h^{[n]} \pi_{n-1}^{[n]} = \varphi^{[n]} U_{\lambda h}^{[n]} \pi_{n-1}^{[n]} = \varphi^{[n]} \Pi_{n-1}^{[n]} U_{\lambda h}^{[n-1]} = \varphi^{[n-1]} U_{\lambda h}^{[n-1]} = h^{[n-1]}.$$

Since $U(V, c, b)$ is the direct limit of the direct system

$$U_V^{[0]} \xrightarrow{\pi_0^1} U_V^{[1]} \xrightarrow{\pi_1^2} U_V^{[2]} \xrightarrow{\pi_2^3} \dots$$

of braided bialgebras, we get that the compatible family $(h^{[n]} : U_V^{[n]} \rightarrow A)_{n \in \mathbb{N}}$ induces a braided bialgebra homomorphism $\bar{h} : U(V, c, b) \rightarrow A$ such that $\bar{h} \pi_n^\infty = h^{[n]}$. In particular

$$\bar{h} i_U = \bar{h} \pi_0^\infty i_V^{[0]} = h^{[0]} i_V^{[0]} = \varphi^{[0]} U_{\lambda h}^{[0]} i_V^{[0]} = \varphi^{[0]} i_A^{[0]} \lambda h = \lambda h.$$

$\square$

The following Theorem 6.8 and Theorem 6.9 constitutes together a Milnor-Moore type theorem for primitively generated braided bialgebras, compare with [MM, Theorem 5.18].

Theorem 6.8. *Let (V, c, b) be a braided Lie algebra. Then the infinitesimal braided Lie algebra of $U(V, c, b)$ is (V, c, b) .*

Proof. Let $U := U(V, c, b)$ and let (W, c_W, b_W) be its infinitesimal braided Lie algebra. By Corollary 5.6, the canonical map $i_U : V \rightarrow U$ induces an isomorphism of braided vector spaces between V and W namely the map $h := i^{[\infty]} : V \rightarrow W$. We only have to check this is a morphism of braided brackets. Let us prove that (6.2) holds by complete induction on $n \in \mathbb{N}$. Let $\varphi^{[n]} : U_U^{[n]} \rightarrow U$ be defined as in the proof of Proposition 6.1. For $n = 0$ and $x \in P_V^{[0]}$, we have

$$b_W^{[0]} P_h^{[0]}(x) = b_U^{[0]} P_h^{[0]}(x) = \varphi^{[0]} P_h^{[0]}(x) = \varphi^{[0]} U_h^{[0]}(x) = \pi_0^\infty(x) = \pi_0^\infty i^{[0]} b^{[0]}(x) = h b^{[0]}(x)$$

so that $b_W^{[0]}P_h^{[0]} = hb^{[0]}$. Let $n > 0$ and assume the statement true for all $0 \leq i \leq n-1$. Then, following 6.2, we know that $P_h^{[n]}$ exists. For $x \in P_V^{[n]}$, we have

$$b_W^{[n]}P_h^{[n]}(x) = b_U^{[n]}P_h^{[n]}(x) = \varphi^{[n]}P_h^{[n]}(x) = \varphi^{[n]}U_h^{[n]}(x) = \pi_n^\infty(x) = \pi_n^\infty i^{[n]}b^{[n]}(x) = hb^{[n]}(x)$$

so that $b_W^{[n]}P_h^{[n]} = hb^{[n]}$. $\square$

What follows is the main result of the paper.

Theorem 6.9. *Every primitively generated braided bialgebra is isomorphic as a braided bialgebra to the universal enveloping algebra of its infinitesimal braided Lie algebra.*

Proof. Let A be a primitively generated braided bialgebra with infinitesimal braided Lie algebra (P, c_P, b_P) and set $U := U(P, c_P, b_P)$. By Theorem 6.7, the identity map of P can be lifted to a morphism of braided bialgebras $\varphi : U \rightarrow A$. Since P generates A as a K -algebra, φ is surjective. On the other hand, by Corollary 5.6, $i_U : P \rightarrow U$ induces an isomorphism between P and $P(U)$. Hence the restriction of φ to $P(U)$ is injective. By [Mo, Lemma 5.3.3] φ is injective too. $\square$

Example 6.10. Assume $\text{char}K = 0$. Let V be a vector space. Regard it as a braided vector space through the canonical flip map $c : V \otimes V \rightarrow V \otimes V : x \otimes y \mapsto y \otimes x$. Then the braiding c is of Hecke type with regular mark 1 (regularity, when the mark is 1, just means $\text{char}K = 0$). Thus, by [Ar1, Theorem 6.14], we have that (V, c) is in $\mathcal{S}$.

Let b such that (V, c, b) is a braided Lie algebra. By Theorem 5.10, we have that $U(V, c, b) = U(V, c, b^{[0]}) = \mathbb{U}(V, c, \beta)$, where $\beta : E(V, c) \rightarrow V$ is the restriction of $b^{[0]}$ to $E(V, c)$. By Lemma 5.11, we have that, for all $t \geq 1$, $b^{[t]} = b^{[1]}$. Moreover $b^{[1]}$ is bijective and it is the inverse of $i^{[1]} : V \rightarrow P^{[1]}$. By [Ar2, Theorem 6.3]

$$\mathbb{U}(V, c, \beta) = \frac{T(V, c)}{(x \otimes y - c(x \otimes y) - [x, y] \mid x, y \in V)} = \frac{T(V)}{(x \otimes y - y \otimes x - [x, y] \mid x, y \in V)}$$

where $[-, -] : V \times V \rightarrow V$ is defined by $[x, y] := \beta(x \otimes y - c(x \otimes y)) = b^{[0]}(x \otimes y - y \otimes x)$ for all $x, y \in V$. It is remarkable that $(V, [-, -])$ is an ordinary Lie algebra (see [Ar2, Remark 6.4]) so that $U(V, c, b)$ is nothing but the ordinary universal enveloping algebra of $(V, [-, -])$.

Conversely, let $[-, -] : V \times V \rightarrow V$ be such that $(V, [-, -])$ is an ordinary Lie algebra. Let A be the ordinary universal enveloping algebra of $(V, [-, -])$. This is an ordinary Hopf algebra, see e.g. [Ka, Proposition V.2.4, page 97] or [Bo, Proposition 7, page 115]. In particular it is a braided bialgebra with braiding the canonical flip map on A . It is primitively generated as, by construction it is generated by the image of V in A . By Theorem 6.9, $A \cong U(P, c_P, b_P)$ where (P, c_P, b_P) is the infinitesimal braided Lie algebra of A . Now, since $\text{char}K = 0$, we have that the canonical map $\sigma : V \rightarrow P = P(A)$ is bijective [Bo, Corollary at page 117]. Since c_P is the restriction of the braiding of A , then c_P is the canonical flip map c on V . Hence we can identify (P, c_P) with (V, c) and σ with the identity of V . Hence we get a braided bialgebra (V, c, b) where $b = b_P$. By construction the first part we have that, for all $t \geq 1$, $b^{[t]} = b^{[1]}$. Moreover $b^{[1]}$ is the inverse of σ so that $b^{[1]} = \text{Id}_V$. On the other hand, one has $b^{[0]}(z) = \varphi^{[0]}(z)$ for all $P(T)$ where $T := T(V, c)$ and $\varphi^{[0]} : U^{[0]} \rightarrow A$ is defined as in the proof of Proposition 6.1. Explicitly, if $z = \sum_{0 \leq i \leq n} z_i$ for some $z_i \in V^{\otimes i}$, then $b^{[0]}(z) = \sum_{0 \leq i \leq n} \varphi^{[0]}(z_i) = \sum_{0 \leq i \leq n} m_A^{i-1}(z_i)$ where $m_A^{i-1} : A^{\otimes i} \rightarrow A : x_1 \otimes \cdots \otimes x_i \mapsto x_1 \cdots x_i$ denotes the iterated multiplication of A .

We are now able to recover the classical Milnor-Moore theorem.

Corollary 6.11. (cf. [MM, Theorem 5.18]) *Let A be a cocommutative ordinary bialgebra in characteristic zero. Then A is isomorphic as a bialgebra to the universal enveloping algebra of $P(A)$.*

Proof. In vie of Example 2.8, A is primitively generated. By Theorem 6.9, A is isomorphic as a braided bialgebra to the universal enveloping algebra of its infinitesimal braided Lie algebra (P, c_P, b_P) . By Example 6.10, $U(P, c_P, b_P)$ is the ordinary universal enveloping algebra of the ordinary Lie algebra $(P, [-, -])$, where $[x, y] := b_P^{[0]}(x \otimes y - y \otimes x) = x \cdot_A y - y \cdot_A x$, for all $x, y \in P$. $\square$

6.12. Let A be a braided bialgebra with infinitesimal braided Lie algebra (V, c, b) . Let $h = \text{Id}_V$. and let $U := U(V, c, b)$. Let $h^{[n]} : U^{[n]} \rightarrow A$ be defined as in the proof of Theorem 6.7. Mimic the construction of [Kh1, Section 3] (see also [Mas, Section 3]) as follows. Let $I := \text{Ker}(h^{[0]})$. Set $I_0 := 0$, $I_1 := (P[I])$, $I_2/I_1 := (P[I/I_1])$, $I_3/I_2 := (P[I/I_2])$ and so on. Continuing this way one gets an increasing chain of ideals of $U^{[0]} = T(V, c)$

$$0 = I_0 \leq I_1 \leq I_2 \leq \cdots \leq I_n \leq \cdots$$

Proposition 6.13. *Keep the assumptions and notations of 6.12. Then, for all $n \in \mathbb{N}$*

$$\text{Ker}(h^{[n]}) = \pi_0^n(I) \quad \text{and} \quad \text{Ker}(\pi_0^n) = I_n,$$

where $\pi_0^n : T(V, c) = U^{[0]} \rightarrow U^{[n]}$ is the canonical projection. Moreover

$$\text{Ker}(\pi_0^\infty) = \bigcup_{n \in \mathbb{N}} I_n.$$

In particular $I = \bigcup_{n \in \mathbb{N}} I_n$ whenever A is primitively generated.

Proof. We proceed by induction on $n \in \mathbb{N}$. For $n = 0$, there is nothing to prove. Let $n > 1$ and assume the formulas true for $n - 1$. Then, since $U^{[n]} = U^{[n-1]}/\text{Ker}(\pi_{n-1}^n)$, we get

$$\text{Ker}(h^{[n]}) = \frac{\text{Ker}(h^{[n-1]})}{\text{Ker}(\pi_{n-1}^n)} = \frac{\pi_0^{n-1}(I)}{\text{Ker}(\pi_{n-1}^n)} = \pi_{n-1}^n \pi_0^{n-1}(I) = \pi_0^n(I).$$

Let us check that $I_n = \text{Ker}(\pi_0^n)$. We have

$$P[\pi_0^{n-1}(I)] = P(\text{Ker}(h^{[n-1]})) = P^{[n-1]} \cap \text{Ker}(h^{[n-1]}) = \text{Ker}(h_{P^{[n-1]}}^{[n-1]}) = \text{Ker}[b^{[n-1]}].$$

Thus, by Corollary 5.8,

$$(6.4) \quad \text{Ker}(\pi_{n-1}^n) = (\text{Ker}[b^{[n-1]}]) = (P[\pi_0^{n-1}(I)]).$$

Now, since $I_{n-1} = \text{Ker}(\pi_0^{n-1})$, then the map

$$\alpha_{n-1} : U^{[0]}/I_{n-1} \rightarrow U^{[n-1]} : z + I_{n-1} \mapsto \pi_0^{n-1}(z)$$

is a well defined braided bialgebra isomorphism. Thus

$$I_n = \text{Ker}(\pi_0^n) \Leftrightarrow \frac{I_n}{I_{n-1}} = \frac{\text{Ker}(\pi_0^n)}{I_{n-1}} \Leftrightarrow \alpha_{n-1} \left[\frac{I_n}{I_{n-1}} \right] = \alpha_{n-1} \left[\frac{\text{Ker}(\pi_0^n)}{I_{n-1}} \right].$$

Since α_{n-1} is a braided bialgebra isomorphism, we get

$$\begin{aligned} \alpha_{n-1} \left[\frac{I_n}{I_{n-1}} \right] &= \alpha_{n-1} \left[\left(P \left[\frac{I}{I_{n-1}} \right] \right) \right] = \left(\alpha_{n-1} \left[P \left[\frac{I}{I_{n-1}} \right] \right] \right) \\ &= \left(P \left[\alpha_{n-1} \left[\frac{I}{I_{n-1}} \right] \right] \right) = (P[\pi_0^{n-1}(I)]) \end{aligned}$$

and

$$\begin{aligned} \alpha_{n-1} \left[\frac{\text{Ker}(\pi_0^n)}{I_{n-1}} \right] &= \pi_0^{n-1}[\text{Ker}(\pi_0^n)] = \pi_0^{n-1}[\text{Ker}(\pi_{n-1}^n \pi_0^{n-1})] \\ &= \pi_0^{n-1}(\pi_0^{n-1})^{-1}[\text{Ker}(\pi_{n-1}^n)] = \text{Ker}(\pi_{n-1}^n) \\ &\stackrel{(6.4)}{=} (P[\pi_0^{n-1}(I)]) \end{aligned}$$

Hence $I_n = \text{Ker}(\pi_0^n)$. Now, we have

$$\text{Ker}(\pi_0^\infty) = \text{Ker}(\varinjlim \pi_0^n) = \sum_{n \in \mathbb{N}} \text{Ker}(\pi_0^n) = \sum_{n \in \mathbb{N}} I_n = \bigcup_{n \in \mathbb{N}} I_n.$$

By Theorem 6.9, if A is primitively generated then there is an isomorphism $h^{[\infty]} : U \rightarrow A$ such that $h^{[\infty]} \circ \pi_0^\infty = h^{[0]}$. Thus $\text{Ker}(h^{[0]}) = \text{Ker}(\pi_0^\infty)$ or, equivalently, one has $I = \bigcup_{n \in \mathbb{N}} I_n$. $\square$

7. THE TRIVIAL BRACKET

In this section we associate a bracket to any braided vector space. We give a condition guaranteeing that the bracket of a given braided Lie algebra is trivial. Finally we check this condition holds on a specific example.

Example 7.1 (The triivial bracket.). We now give a construction of a trivial braided Lie algebra structure for any braided vector space (V, c) . Let $(U^{[0]}, i^{[0]})$ be defined as in Definition 5.1. Let $b_{tr}^{[0]} : P^{[0]} = P(U^{[0]}) \rightarrow V$ be the restriction of the canonical projection $U^{[0]} \rightarrow V$. Since this projection is a morphism of braided vector spaces, it is clear that $b_{tr}^{[0]}$ fulfills (5.1) and (5.2) for $n = 0$. As in Definition 5.1 we can build $(U^{[1]}, i^{[1]})$. Since $b_{tr}^{[0]} i^{[0]} = \text{Id}_V$ we get

$$U^{[1]} = \frac{U^{[0]}}{\left(\left[\text{Id} - i^{[0]} b_{tr}^{[0]} \right] [P^{[0]}] \right)} = \frac{U^{[0]}}{\left(\text{Ker} \left[b_{tr}^{[0]} \right] \right)}.$$

Note that $\text{Ker} \left[b_{tr}^{[0]} \right]$ is the space spanned by primitive elements in $T(V, c)$ of homogeneous degree at least 2. Now $U^{[1]}$ is a graded braided bialgebra. The component of degree one is $V^{[1]} = \text{Im}(i^{[1]}) \cong V$ so that we can consider the canonical projection $U^{[1]} \rightarrow V$. Moreover, its restriction $b_{tr}^{[1]} : P^{[1]} = P(U^{[1]}) \rightarrow V$ fulfills (5.1) and (5.2) for $n = 1$. Proceeding this way one gets a bracket $b_{tr} = \left(b_{tr}^{[n]} \right)_{n \in \mathbb{N}}$ on the braided vector space (V, c) that will be called the **trivial bracket** of (V, c) . By Proposition 5.7, (V, c, b_{tr}) is a braided Lie algebra. The universal enveloping algebra of rank n in this case will be also denoted by

$$S^{[n]} := S^{[n]}(V, c) := U \left(V, c, \left(b_{tr}^{[i]} \right)_{0 \leq i \leq n-1} \right)$$

and it will be called the **symmetric algebra of rank n of (V, c)** . It is the braided bialgebra $S^{[n]}(B)$ introduced in [Ar1, Definition 3.11] in the case $B = U^{[0]}$. Thus, since the tensor algebra $U^{[0]}$ is strongly $\mathbb{N}$ -graded as an algebra, by applying [Ar1, Corollary A.11], we obtain that

$$U(V, c, b_r) = \varinjlim U^{[n]} \left(V, c, \left(b_{tr}^{[i]} \right)_{0 \leq i \leq n-1} \right) = \varinjlim S^{[n]} = S^{[\infty]}$$

identifies with the Nichols algebra $\mathcal{B}(V, c)$.

Proposition 7.2. *Let (V, c, b) be a braided Lie algebra. Let $p_0^\infty : T(V, c) \rightarrow \mathcal{B}(V, c)$ and $\pi_0^\infty : T(V, c) \rightarrow U(V, c, b)$ be the canonical projections. Suppose the ideal $\text{Ker}(p_0^\infty)$ is generated by a set W . The following are equivalent.*

- (i) π_0^∞ vanishes on W .
- (ii) b is the trivial bracket of (V, c) .

Proof. (i) $\Rightarrow$ (ii) Since $\text{Ker}(p_0^\infty) = (W)$ and π_0^∞ vanishes on W , there exists a unique algebra homomorphism $\phi : \mathcal{B}(V, c) \rightarrow U(V, c, b)$ such that $\phi \circ p_0^\infty = \pi_0^\infty$. From the surjectivity of p_0^∞ one deduce that ϕ is indeed a braided bialgebra homomorphism. Moreover ϕ is surjective as π_0^∞ is. Note that

$$\phi|_V = (\phi \circ p_0^\infty)|_V = (\pi_0^\infty)|_V = i_U$$

which is injective. Since $P(\mathcal{B}(V, c)) \cong V$, by [Mo, Lemma 5.3.3], ϕ is injective whence bijective.

This isomorphism yields an isomorphism of braided brackets between the infinitesimal part of $\mathcal{B}(V, c) = U(V, c, b_r)$ and the infinitesimal part of $U(V, c, b)$. In view of Theorem 6.8, we get that there is an isomorphism of braided brackets between $h : (V, c, b_r) \rightarrow (V, c, b)$. By definition of ϕ it is also clear that $h = \text{Id}_V$. By (6.2), for each $n \in \mathbb{N}$, we get $b_r^{[n]} = h b_r^{[n]} = b^{[n]} P_h^{[n]} = b^{[n]}$ so that $b = b_r$.

(ii) $\Rightarrow$ (i) Since b is the trivial bracket of (V, c) , by Example 7.1, we have that $U(V, c, b) = \mathcal{B}(V, c)$ and $\pi_0^\infty = p_0^\infty$. $\square$

We now give an example of a braided vector space (V, c) which becomes a braided Lie algebra only in a trivial way. It is remarkable that (V, c) does not belong to the class $\mathcal{S}$ of Remark 5.9. Indeed this gives a counterexample to Theorem 5.10. In fact

$$U(V, c, b_{tr}) = \mathcal{B}(V, c) \neq S^{[1]}(V, c) = U(V, c, b_{tr}^{[0]}).$$

Example 7.3. Let K be a fixed field with $\text{char}(K) \neq 2$. At the end of [Kh3], an example of a two dimensional braided vector space (V, c) of combinatorial rank 2 is given. This essentially means that $S^{[0]}(V, c) \neq S^{[1]}(V, c) \neq S^{[2]}(V, c) = \mathcal{B}(V, c)$. The braiding c is of diagonal type of the form

$$c(x_i \otimes x_j) = q_{i,j} x_j \otimes x_i, 1 \leq i, j \leq 2.$$

where x_1, x_2 is a basis of V over K , $q_{1,2} = 1 \neq -1$ and $q_{i,j} = -1$ for all $(i, j) \neq (1, 2)$. One has that

$$\mathcal{B}(V, c) = \frac{T(V, c)}{(x_1^2, \quad x_2^2, \quad x_1 x_2 x_1 x_2 + x_2 x_1 x_2 x_1)}.$$

Thus $\mathcal{B}(V, c)$ has basis

$$1_K, x_1, x_2, x_2 x_1, x_1 x_2, x_1 x_2 x_1, x_2 x_1 x_2, x_2 x_1 x_2 x_1.$$

Note that the endomorphism $c : V \otimes V \rightarrow V \otimes V$ has minimal polynomial $(X + 1)(X^2 + 1)$. Our aim here is to determine all primitively generated connected braided bialgebras A with infinitesimal part (V, c) . Let A be such a braided bialgebra. By Theorem 6.9, A is isomorphic to the enveloping algebra $U(P, c_P, b_P)$ associated to its infinitesimal braided Lie algebra (P, c_P, b_P) . Therefore we have to investigate the possible braided Lie algebra structures on (V, c) .

Let b be a bracket on (V, c) such that (V, c, b) is a braided Lie algebra. Set $T := T(V, c)$. Clearly $x_1^2, x_2^2 \in P^{[0]} = P(T)$.

Suppose there is $t \in \{1, 2\}$ such that $b^{[0]}(x_t^2) \neq 0$. Note that x_t^2 is a homogeneous quantum operation in the sense of [Ar2, Definition 9.1]. By [Ar2, Corollary 9.6], there is $a \in \{1, 2\}$ such that the following relation holds

$$q_{a,j} = \prod_{l=1,2} q_{l,j}^{d_l} = q_{t,j}^2 = 1, \text{ for all } j = 1, 2.$$

In particular $-1 = q_{a,1} = 1$ which contradicts the hypothesis $\text{char}(K) \neq 2$. Hence we proved that

$$b^{[0]}(x_t^2) = 0 \text{ for } t = 1, 2.$$

Therefore

$$(7.1) \quad \pi_0^1(x_t^2) = \pi_0^1[i^{[0]}b^{[0]}(x_t^2)] = 0, \text{ for } t = 1, 2.$$

Using (7.1), one easily checks that $w := \pi_0^1(x_1 x_2 x_1 x_2 + x_2 x_1 x_2 x_1) \in P^{[1]} = P(U^{[1]})$.

Let us check that $b^{[1]}[w] = 0$. Set $b^{[1]}(w) = b_1^{[1]}x_1 + b_2^{[1]}x_2$. By (5.1), for $1 \leq j \leq n$ we have that

$$c_{V^{[1]}}(i^{[1]}b^{[1]} \otimes V^{[1]}) = (V^{[1]} \otimes i^{[1]}b^{[1]})c_{P^{[1]}, V^{[1]}},$$

Now

$$\begin{aligned} c_{V^{[1]}}(i^{[1]}b^{[1]} \otimes V^{[1]})(w \otimes \pi_0^1(x_j)) &= c_{V^{[1]}}(i^{[1]}b^{[1]}(w) \otimes \pi_0^1(x_j)) \\ &= b_1^{[1]}c_{V^{[1]}}(\pi_0^1(x_1) \otimes \pi_0^1(x_j)) + b_2^{[1]}c_{V^{[1]}}(\pi_0^1(x_2) \otimes \pi_0^1(x_j)) \\ &= (\pi_0^1 \otimes \pi_0^1)[b_1^{[1]}c(x_1 \otimes x_j) + b_2^{[1]}c(x_2 \otimes x_j)] \\ &= q_{1,j}b_1^{[1]}\pi_0^1(x_j) \otimes \pi_0^1(x_1) + q_{2,j}b_2^{[1]}\pi_0^1(x_j) \otimes \pi_0^1(x_2) \\ &= q_{1,j}b_1^{[1]}\pi_0^1(x_j) \otimes \pi_0^1(x_1) - b_2^{[1]}\pi_0^1(x_j) \otimes \pi_0^1(x_2) \end{aligned}$$

and

$$\begin{aligned} (V^{[1]} \otimes i^{[1]}b^{[1]})c_{P^{[1]}, V^{[1]}}(w \otimes \pi_0^1(x_j)) &= q_{1,j}^2 q_{2,j}^2 (V^{[1]} \otimes i^{[1]}b^{[1]})(\pi_0^1(x_j) \otimes w) \\ &= b_1^{[1]}\pi_0^1(x_j) \otimes \pi_0^1(x_1) + b_2^{[1]}\pi_0^1(x_j) \otimes \pi_0^1(x_2). \end{aligned}$$

Since $\pi_0^1(x_i) = i^{[0]}(x_j)$ and $i^{[0]}$ is injective, we deduce that

$$q_{1,j}b_1^{[1]}x_j \otimes x_1 - b_2^{[1]}x_j \otimes x_2 = b_1^{[1]}x_j \otimes x_1 + b_2^{[1]}x_j \otimes x_2$$

whence, for $j = 1$ we get

$$2b_1^{[1]} = 0 = 2b_2^{[1]}$$

whence $b^{[1]}[w] = 0$ as $\text{char}(K) \neq 2$. Now

$$\begin{aligned} \pi_0^\infty(x_1x_2x_1x_2 + x_2x_1x_2x_1) &= \pi_1^\infty\pi_0^1(x_1x_2x_1x_2 + x_2x_1x_2x_1) \\ &= \pi_1^\infty(w) \\ &= \pi_2^\infty\pi_1^2(w) \\ &\stackrel{w \in P^{[1]}}{=} \pi_2^\infty\pi_1^2i^{[1]}b^{[1]}(w) = 0. \end{aligned}$$

Since we already proved that $\pi_0^1(x_t^2) = 0$, we also have $\pi_0^\infty(x_t^2) = 0$ for $t = 1, 2$. By Proposition 7.2, b is the trivial bracket of (V, c) . As a consequence, we get $U(V, c, b) = \mathcal{B}(V, c)$. In view of Theorem 6.9, $\mathcal{B}(V, c)$ is then the unique primitively generated connected braided bialgebra A with infinitesimal part (V, c) .

REFERENCES

- [AN] N. Andruskiewitsch, S. Natale, *Braided Hopf algebras arising from matched pairs of groups*. J. Pure Appl. Algebra **182** (2003), no. 2-3, 119–149.
- [AS1] N. Andruskiewitsch, H.-J. Schneider, *Lifting of quantum linear spaces and pointed Hopf algebras of order p^3* . J. Algebra **209** (1998), no. 2, 658–691.
- [AS2] N. Andruskiewitsch, H.-J. Schneider, *Finite quantum groups and Cartan matrices*. Adv. Math. **154** (2000), no. 1, 1–45.
- [AS3] N. Andruskiewitsch, H.-J. Schneider, *Pointed Hopf algebras*. New directions in Hopf algebras, 1–68, Math. Sci. Res. Inst. Publ., **43**, Cambridge Univ. Press, Cambridge, 2002.
- [AMS] A. Ardizzoni, C. Menini and D. Ştefan, *Braided Bialgebras of Hecke-type*, J. Algebra, Vol. **321** (2009), 847–865.
- [Ar1] A. Ardizzoni, *On the Strongness Degree of a Graded Braided Bialgebra*, submitted. (<http://web.unife.it/utenti/alessandro.ardizzoni/Pdf/Sdeg.pdf>)
- [Ar2] A. Ardizzoni, *On Primitively Generated Braided Bialgebras*, submitted. (<http://web.unife.it/utenti/alessandro.ardizzoni/Pdf/OnPrimGen.pdf>)
- [Ba] J.C. Baez, *Hochschild homology in a braided tensor category*, Trans. Amer. Math. Soc. **334** (1994), 885–906.
- [Bo] N. Bourbaki, *Lie groups and Lie algebras. Chapters 1–3*. Translated from the French. Reprint of the 1989 English translation. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 1998.
- [Ka] C. Kassel, *Quantum groups*. Graduate Texts in Mathematics, 155. Springer-Verlag, New York, 1995.
- [Kh1] Kharchenko, V. K. A combinatorial approach to the quantification of Lie algebras. (English summary) Pacific J. Math. **203** (2002), no. 1, 191–233.
- [Kh2] V. K. Kharchenko, *Connected Braided Hopf algebras*, J. Alg. **307** (2007), 24–48.
- [Kh3] V. K. Kharchenko, *Skew primitive elements in Hopf algebras and related identities*. J. Algebra **238** (2001), no. 2, 534–559.
- [Mas] A. Masuoka, *Construction of quantized enveloping algebras by cocycle deformation*. (English, Arabic summary) Arab. J. Sci. Eng. Sect. C Theme Issues **33** (2008), no. 2, 387–406.
- [MM] J. W. Milnor and J. C. Moore, *On the structure of Hopf algebras*. Ann. of Math. (2) **81** (1965) 211–264.
- [Mo] S. Montgomery, *Hopf Algebras and their actions on rings*, CMBS Regional Conference Series in Mathematics **82** (1993).
- [Sw] M. Sweedler, *Hopf Algebras*, Benjamin, New York, 1969.
- [Ta] M. Takeuchi, *Survey of braided Hopf algebras*, Contemp. Math. **267** (2000), 301–324.

Current address: University of Ferrara, Department of Mathematics, Via Machiavelli 35, Ferrara, I-44121, Italy

E-mail address: alessandro.ardizzoni@unife.it

URL: <http://www.unife.it/utenti/alessandro.ardizzoni>